



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>A61K 38/17</b>	<b>A2</b>	(11) International Publication Number: <b>WO 00/37094</b> (43) International Publication Date: 29 June 2000 (29.06.00)								
<p>(21) International Application Number: PCT/US99/30734</p> <p>(22) International Filing Date: 21 December 1999 (21.12.99)</p> <p>(30) Priority Data:</p> <table border="0"> <tr> <td>60/113,407</td> <td>22 December 1998 (22.12.98)</td> <td>US</td> </tr> <tr> <td>PCT/US99/06797</td> <td>30 March 1999 (30.03.99)</td> <td>US</td> </tr> <tr> <td>60/172,239</td> <td>20 October 1999 (20.10.99)</td> <td>US</td> </tr> </table> <p>(71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): COHEN, Fredric, Jay [US/US]; 8330 Catamaran Drive, Indianapolis, IN 46236 (US). POSADA, James, Arthur [US/US]; 3575 Inverness Boulevard, Carmel, IN 46032 (US). WIERDA, Daniel [US/US]; 2606 South Brandywine Trail, Greenfield, IN 46140 (US).</p> <p>(74) Agents: WEBSTER, Thomas, D. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).</p>	60/113,407	22 December 1998 (22.12.98)	US	PCT/US99/06797	30 March 1999 (30.03.99)	US	60/172,239	20 October 1999 (20.10.99)	US	<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> Without international search report and to be republished upon receipt of that report.</p>
60/113,407	22 December 1998 (22.12.98)	US								
PCT/US99/06797	30 March 1999 (30.03.99)	US								
60/172,239	20 October 1999 (20.10.99)	US								
<p>(54) Title: THERAPEUTIC APPLICATIONS OF FLINT POLYPEPTIDES</p> <p>(57) Abstract</p> <p>Mature FLINT protein (mFLINT) binds FasL and LIGHT, and prevents FasL-Fas interaction. mFLINT inhibits FasL-Fas-mediated apoptotic and proinflammatory activity, and is useful in treating disorders that may be associated with abnormal apoptosis and inflammation. The invention provides the amino acid and nucleotide sequences of FLINT and mature FLINT. Therapeutic compositions and methods of treatment utilizing mFLINT also are provided.</p>										

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-1-

## THERAPEUTIC APPLICATIONS OF FLINT POLYPEPTIDES

CROSS REFERENCE

5        This application claims the benefit of U.S. Provisional Application No. 60/113,407, filed December 22, 1998; PCT Application No. PCT US99/06797, filed March 30, 1999; and U.S. Provisional Application filed October 20, 1999.

BACKGROUND OF THE INVENTION

10        FasL (also called CD95L and APO1L) is expressed on various cell types and can produce biological responses such as proliferation, differentiation, immunoregulation, inflammatory response, cytotoxicity, and apoptosis.

15        Interestingly, mutations in FasL, the ligand for the TNFR-family receptor FAS/APO (Suda et al., 1993, Cell 75:1169-78, are associated with autoimmunity (Fisher et al., 1995, Cell 81:935-46), while overproduction of FasL may be implicated in drug-induced hepatitis. FasL is expressed in immune-

20        privileged tissues of the eye, testis, brain and some tumors. It has also been found in kidney and lung as well as in activated thymocytes, splenocytes, and T lymphocytes.

      Apoptosis plays a central role in both development and in homeostasis. Cells die by apoptosis in the developing

25        embryo during morphogenesis or synaptogenesis and in the adult animal during tissue turnover or at the end of an immune response. Because the physiological role of apoptosis is crucial, aberration of this process can be detrimental. For example, unscheduled apoptosis of certain

30        brain neurons contributes to disorders such as Alzheimer's and Parkinson's disease, whereas the failure of dividing

-2-

cells to initiate apoptosis after sustaining severe DNA damage contributes to cancer.

Survival signals from the cell's environment and internal sensors for cellular integrity normally keep a cell's apoptotic machinery in check. In the event that a cell loses contact with its surroundings or sustains irreparable damage, the cell initiates apoptosis. A cell that simultaneously receives conflicting signals driving or attenuating its division cycle also triggers apoptosis.

10 Mammals have evolved yet another mechanism that enables the organism actively to direct individual cells to self-destruct. This kind of "instructive" apoptosis is important especially in the immune system. Death receptors-cell surface receptors that transmit apoptosis signals initiated

15 by specific "death ligands" - play a central role in instructive role apoptosis. These receptors can activate death caspases within seconds of ligands binding, causing an apoptotic demise of the cell within hours.

Death receptors belong to the tumor necrosis factor (TNF) receptor gene superfamily, which is defined by similar, cysteine-rich extracellular domains. The death receptors contain in addition a homologous cytoplasmic sequence termed the "death domain". Death domains typically enable death receptors to engage the cell's apoptotic

20 machinery, but in some instances they mediate functions that are distinct from or even counteract apoptosis.

Fas (also called CD95 or Apol) is a well-characterized death receptor. Fas and Fas ligand (FasL) play an important role in apoptosis. Fas L is a homotrimeric

30 molecule. It is suggested that each FasL trimer binds three Fas molecules. Because death domains have a propensity to

-3-

associate with one another, Fas ligation leads to clustering of the receptors' death domains. An adapter protein called FADD (Fas-associated death domain; also called Mort1) then binds through its own death domain to the clustered receptor death domains. FADD also contains a "death effector domain" that binds to an analogous domain repeated in tandem within the zymogen form of caspase-8 (also called FLICE, or MACH). Upon recruitment by FADD, caspase-8 oligomerization drives its activation through self-cleavage. Caspase-8 then activates downstream effector caspases such as caspase-9 committing the cell to apoptosis. Ashkenazi A., et al. "Death Receptors: Signaling and Modulation Science 281, 1305-1308 (August 1998).

Although it triggers apoptosis in T lymphocytes, FasL is also proinflammatory. FasL has been shown to stimulate neutrophils, also called polymorphonuclear leukocytes (PMNs), activation. (Chen J. et al., Science 282: 1714-17 (1998)) FasL-Fas binding has been implicated in clonal deletions of autoreactive lymphocytes in peripheral lymphoid tissues and in elimination of autoreactive lymphocyte populations, thus contributing to homeostasis of the immune system. However, it has been found that expression of FasL on myotubes or pancreatic islets of transgenic mice induces a granulocytic response that accelerates graft rejection (Allison J. et al., Proc. Natl. Acad. Sci, 94:3943-47 (April 1997); Kang S-M. et al., Nature Medicine, Vol. 3, No. 7, 738-743 (July 1997)).

At least one of the effects of FasL-Fas receptor binding is apoptosis, which is necessary for homeostasis. However, sometimes the balance of ligand-receptors binding is upset in stress, disease or trauma. One of the negative

-4-

effects of unregulated FasL-Fas binding is runaway or aberrant apoptosis. Another effect of said binding is the destruction of healthy cells caused by neutrophils that have been activated by FasL.

5 For example, one of the more tragic outcomes of runaway apoptosis in a specific organ is acute liver failure. Acute liver failure is characterized by an over-activation of the apoptotic pathway where there is massive apoptosis of hepatocytes and hemorrhagic liver changes. Acute liver  
10 failure can happen as a result of viral infections affecting the liver, bacterial infections affecting the liver, hepatitis, hepatocellular injury and/or other conditions where hepatocytes undergo massive apoptosis. One example of an infection resulting in acute liver failure is bacterium-  
15 induced fulminant hepatitis.

#### SUMMARY OF THE INVENTION

The invention encompasses an isolated nucleic acid molecule having the sequence of Figure 1, an isolated  
20 nucleic acid molecule having the sequence of Figure 3, isolated polypeptide having the sequence of Figure 1, and an isolated polypeptide having the sequence of Figure 3. The invention also includes a transgenic mouse comprising a transgene having the sequence of Figure 1.

25 According to one aspect of the invention, mFLINT is provided for use in treating the following conditions: acute liver failure; inflammation of the liver; abnormal hepatocyte apoptosis; sepsis; a disorder associated with inflammation; hepatitis; treating abnormal apoptosis; an  
30 ischemia-associated injury or disorder; injury or disorder is associated with hypercoagulation, including use with

-5-

agent selected from the group selected from thrombolytic and antithrombotic agents, like activated protein C; a reperfusion-associated injury or disorder; damage to a cardiac myocyte resulting from abnormal myocardial ischemia; Type I diabetes; cancer; damage to an innocent bystander tissue that is induced by a chemotherapeutic agent or therapeutic irradiation, including bone marrow, intestinal epithelium, oral cavity epithelium; injury to hematopoietic progenitor cells that have been exposed to therapeutic radiation or chemotherapy; cell damage from therapeutic irradiation or chemotherapy, including damage to intestinal epithelial cells, hematopoietic progenitor cells, and peripheral blood cells; aplastic anemias; myelodysplastic syndromes; and pancytopenic conditions.

According to another aspect of the invention, mFLINT is provided for use in promoting the growth or differentiation of hematopoietic progenitor cells and promoting the growth or differentiation of CD34+ cells.

Further aspects of the invention include formulations, having mFLINT as an active ingredient that are adapted for the treatment of the following: acute liver failure; inflammation of the liver; abnormal hepatocyte apoptosis; sepsis; a disorder associated with inflammation; hepatitis; treating abnormal apoptosis; an ischemia-associated injury or disorder; injury or disorder is associated with hypercoagulation, including use with agent selected from the group selected from thrombolytic and antithrombotic agents, like activated protein C; a reperfusion-associated injury or disorder; damage to a cardiac myocyte resulting from abnormal myocardial ischemia; Type I diabetes; cancer; damage to an innocent bystander tissue that is induced by a

-6-

chemotherapeutic agent or therapeutic irradiation, including bone marrow, intestinal epithelium, oral cavity epithelium; injury to hematopoietic progenitor cells that have been exposed to therapeutic radiation or chemotherapy; cell  
5 damage from therapeutic irradiation or chemotherapy, including damage to intestinal epithelial cells, hematopoietic progenitor cells, and peripheral blood cells; aplastic anemias; myelodysplastic syndromes; and pancytopenic conditions.

10 According to still another aspect of the invention provides formulations, having mFLINT as an active ingredient that are adapted for promoting the growth or differentiation of hematopoietic progenitor cells and promoting the growth or differentiation of CD34+ cells.

15 Yet other embodiments of the invention include the use of mFLINT in the preparation of medicaments useful in treating the following conditions: acute liver failure; inflammation of the liver; abnormal hepatocyte apoptosis; sepsis; a disorder associated with inflammation; hepatitis;  
20 treating abnormal apoptosis; an ischemia-associated injury or disorder; injury or disorder is associated with hypercoagulation, including use with agent selected from the group selected from thrombolytic and antithrombotic agents, like activated protein C; a reperfusion-associated injury or  
25 disorder; damage to a cardiac myocyte resulting from abnormal myocardial ischemia; Type I diabetes; cancer; damage to an innocent bystander tissue that is induced by a chemotherapeutic agent or therapeutic irradiation, including bone marrow, intestinal epithelium, oral cavity epithelium;  
30 injury to hematopoietic progenitor cells that have been exposed to therapeutic radiation or chemotherapy; cell



-7-

damage from therapeutic irradiation or chemotherapy,  
including damage to intestinal epithelial cells,  
hematopoietic progenitor cells, and peripheral blood cells;  
aplastic anemias; myelodysplastic syndromes; and  
5 pancytopenic conditions.

According to an additional aspect, the invention  
include the use of mFLINT in the preparation of medicaments  
useful in promoting the growth or differentiation of  
hematopoietic progenitor cells and promoting the growth or  
10 differentiation of CD34+ cells.

The present invention encompasses a method for treating  
an individual suffering from abnormal hepatocyte apoptosis,  
comprising administration of a therapeutically amount of  
mFLINT protein to said individual. The invention also  
15 includes a method of treating an individual suffering from a  
disorder associated with inflammation, comprising  
administration of a therapeutically amount of mFLINT protein  
to said individual. Additionally, the invention includes a  
method of treating an individual suffering from abnormal  
20 apoptosis, comprising administration of a therapeutically  
effective amount of mFLINT protein to said individual.

The present invention further includes the use of  
mFLINT to treat a variety of disorders of the liver. In  
this regard, the invention encompasses a method of treating  
25 an individual suffering from acute liver failure, comprising  
administration of a therapeutically amount of mFLINT protein  
to said individual. The invention also includes a method of  
treating an individual suffering from inflammation of the  
liver, comprising administration of a therapeutically amount  
30 of mFLINT protein to said individual. Another method of the  
invention is for treating an individual suffering from

-8-

hepatitis, comprising administration of a therapeutically effective amount of mFLINT protein to said individual.

The invention also encompasses a method of treating an individual suffering from sepsis, comprising administration  
5 of a therapeutically amount of mFLINT protein to said individual.

Also included within the present invention is a method for treating an individual suffering from an ischemia-associated injury or disorder, comprising administration of  
10 a therapeutically effective amount of mFLINT protein to said individual. Such an injury or disorder may associated with hypercoagulation. In this regard, the invention also includes treating a disorder associated with hypercoagulation, comprising administration of a  
15 therapeutically effective amount of mFLINT, in combination with an thrombolytic agent or an antithrombotic agent. One example of such an antithrombotic agent is activated protein C.

The invention also includes a method of treating an  
20 individual suffering from a reperfusion-associated injury or disorder, comprising administration of a therapeutically effective amount of mFLINT protein to said individual.

The invention further encompasses a method of preventing damage to a cardiac myocyte in an individual that  
25 has suffered from abnormal myocardial ischemia, comprising administration of a therapeutically effective amount of mFLINT protein to said individual.

Also included in the invention is a method of treating an individual suffering from Type I diabetes, comprising  
30 administration of a therapeutically amount of mFLINT protein to said individual. Yet another method encompassed by the

-9-

present invention is a method of treating an individual suffering from cancer, comprising administration of a therapeutically effective amount of mFLINT protein to said individual.

5       The invention also includes a method of treating damage to an innocent bystander tissue that is induced by a chemotherapeutic agent or therapeutic irradiation, in an individual treated with said agent or irradiation, comprising administration of a therapeutically effective  
10 amount of mFLINT to said individual. Such tissues include, bone marrow and intestinal epithelium, including the epithelium of the oral cavity.

      The invention encompasses method for treating hematopoietic progenitor cells that have been exposed to  
15 therapeutic radiation or chemotherapy, comprising administering mFLINT to said cells. Such a method promotes the recover of hematopoietic progenitor cells from the adverse effects of therapeutic radiation or chemotherapy. The invention also includes a method of promoting the growth  
20 or differentiation of a hematopoietic progenitor cell, comprising administering mFLINT to said cell. The invention further includes a method of promoting the growth or differentiation of a CD34+ cell, comprising administering mFLINT to said cell.

25       The invention also includes a method for treating cancer, comprising treating bone marrow cells *in vitro* with mFLINT, and administering said cells to said patient, wherein said administration occurs after said patient has been treated with therapeutic irradiation or chemotherapy.  
30 The bone marrow cells may be autologous, i.e., from the

-10-

patient being treated, or heterologous, i.e., from an individual other than the patient.

The invention also encompasses a method of treating cell damage in a patient who receives therapeutic  
5 irradiation or chemotherapy, comprising administering to said patient, a therapeutically effective amount of mFLINT with said irradiation or chemotherapy. The cell damage may be to an intestinal epithelial cell, a hematopoietic progenitor cell, or a peripheral blood cell.

10 The invention also contemplates methods of treating aplastic anemia, myelodysplastic syndrome or a pancytopenic condition, comprising administering a therapeutically effective amount of mFLINT to a patient suffering from aplastic anemia.

15 The invention also relates to a method for treating and/or inhibiting acute respiratory distress syndrome.

The invention also relates to a method for treating and/or inhibiting ulcerative colitis.

The invention also relates to a method for treating  
20 and/or inhibiting ischemia injury during organ transplantation.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the combined amino acid and nucleotide  
25 sequences of human FLINT.

Figure 2 shows the combined amino acid and nucleotide sequences of human FLINT.

Figure 3 shows the combined amino acid and nucleotide sequences of human mFLINT.

30 Figure 4 shows the combined amino acid and nucleotide sequences of human mFLINT.

-11-

Figure 5 compares the performance of mFLINT with other agents in an animal sepsis model after different times of LPS administration.

Figure 6, left panel, shows an experiment comparing mFLINT with anti-TNF in an animal sepsis model. The right panel compares a lower dose of mFLINT against anti-TNF in the same model.

Figure 7 shows an experiment comparing intravenous and intraperitoneal delivery of mFLINT and anti-TNF in an animal sepsis model.

Figure 8 shows reduction in B16 melanoma tumor volume in response to treatment with mFLINT.

Figure 9 shows mFLINT promoted recovery of bone marrow progenitor cells following irradiation.

Figure 10 shows mFLINT promoted recovery of bone marrow progenitor cells following treatment with 5-fluorouracil.

#### DETAILED DESCRIPTION OF THE INVENTION

By employing methods for identifying compounds that bind Fas ligand, applicants have discovered that human FLINT polypeptides are capable of disrupting FasL-Fas receptor interaction. Applicants have discovered methods for modulating the TNFR proteins and their respective ligand interactions where such interactions cause or exacerbate disease and methods for preventing or treating diseases.

As noted above, it is well established that one of the downstream effects of FasL-Fas receptor binding is apoptosis. In conditions evidencing abnormal activation of this pathway, runaway apoptosis results, which is a contributing factor in the pathology of a variety of disorders. Yet another downstream effect of FasL-Fas

-12-

receptor binding is neutrophil activation wherein cells are destroyed by neutrophils activated by FasL.

Recently, it was discovered that FasL induces in peritoneal exudate cells (PEC) the processing and release of IL-1 $\beta$  that is responsible for the neutrophil infiltration. In particular, Miwa K., et al. Nature Medicine, 4(11): 1287-1292 (Nov. 1998), found that inoculation of tumor cells expressing Fas ligand into wild-type mice induces a massive neutrophil infiltration that is, in contrast, suppressed in IL-1 $\alpha\beta$  knockout mice. This indicates that FasL has an inflammatory role. It also suggests that apoptosis may itself induce inflammation under certain conditions. In addition, it is known that certain inflammatory factors can induce the Fas-mediated apoptosis pathway. Thus, it is likely that FasL acts via two possibly distinct yet related pathways in exerting its pathological effects.

The inventors discovered that FLINT polypeptides bind to FasL with at least the same, if not greater, affinity than the Fas receptor itself. As a result of binding FasL, FLINT polypeptides can interfere with FasL binding to Fas receptor and interfere with events downstream. Using various *in vitro* and animal models, presented below in the examples, the inventors have demonstrated the ability of FLINT polypeptides to prevent Fas-mediated deleterious effects. It is apparent from these data that FLINT can act on both the apoptotic and proinflammatory aspects of FasL activity, which implicates its use in several disease and injury states discussed below.

Data presented below show that mFLINT inhibits both FasL apoptosis-inducing activity and proinflammatory activity. By antagonizing FasL, mFLINT polypeptides can

-13-

modulate the destruction of healthy cells caused both by neutrophils activated by FasL and by apoptotic damage mediated directly by FasL-Fas interaction. Accordingly, the present methods of treatment utilizing mFLINT are useful in the treatment and prevention of disorders associated with the direct apoptotic effects of FasL and/or the damage mediated by the proinflammatory effects of FasL, whether or not these represent distinct physiological pathways.

Thus, as characterized generally, the invention relates to methods preventing or treating conditions caused or exacerbated by "abnormal apoptosis," in particular, apoptosis induced by Fas ligand (FasL) and Fas receptor (Fas) binding (also referred to as FasL-Fas binding). This invention also relates to methods of preventing or treating conditions caused by a proinflammatory response, more particularly, a proinflammatory response caused by FasL induced neutrophil activation.

Throughout this application, headings are used for purposes of organizational convenience only, and should not be construed to limit the subject matter described herein.

#### I. Definitions

The following definitions are used in the present application.

As used in this application, the term "FLINT" refers to any full length FLINT polypeptide. Such a full-length polypeptide includes the leader sequence, which is amino acids 1-29 of a FLINT polypeptide. Examples of FLINT include a protein having the amino acid sequence set forth in Figure 1, which is a human FLINT, and Figure 2, which is a human FLINT variant.

-14-

As used in this application, the term "mFLINT" refers to a mature FLINT, i.e., FLINT which does not have a leader (also known as signal) peptide. Examples of mFLINT include a protein having the amino acid sequence set forth in Figure 3, which is the protein of Figure 1 without its leader peptide, and a protein having the amino acid sequence set forth in Figure 4, which is the protein of Figure 2 without amino acids 1-29. Accordingly, an "mFLINT gene" is a nucleic acid that encodes an mFLINT polypeptide. The nucleic acids set forth in Figures 3 and 4 are examples of mFLINT genes according to the present invention.

As used here, with reference to FasL or Fas expression or interaction, and to any resulting apoptosis, the terms "inappropriate" and "abnormal" should be read to include any deviation from normal expression, interaction or apoptosis levels. Such deviations include temporal, quantitative and qualitative abnormalities. FasL or Fas "expression" refers not only transcription, translation and associated events, but also to any process that results in increased the availability of active FasL or Fas, such as transport and cell surface availability/accessibility.

Also as used herein the term "abnormal apoptosis" refers to excessive and/or improper apoptosis. Typically abnormal apoptosis is observed in cells and tissues that have undergone physical, chemical or biological insult. Such insults include, but are not limited to physical injury, viral infection, bacterial infection, ischemia, irradiation, chemotherapy, and the like.

The term "fusion protein" denotes a hybrid protein molecule not found in nature comprising a translational fusion or enzymatic fusion in which two or more different



-15-

proteins or fragments thereof are covalently linked on a single polypeptide chain.

"Host cell" refers to any eucaryotic or prokaryotic cell that is suitable for propagating and/or expressing a cloned gene contained on a vector that is introduced into  
5 said host cell by, for example, transformation or transfection, or the like.

The term "inhibit" or "inhibiting" includes the generally accepted meaning, which includes prohibiting,  
10 preventing, restraining, slowing, stopping, or reversing progression or severity of a disease or condition.

"Isolated nucleic acid compound" refers to any RNA or DNA sequence, however constructed or synthesized, which is locationally distinct from its natural location.

15 The term "plasmid" refers to an extrachromosomal genetic element. The plasmids disclosed herein are commercially available, publicly available on an unrestricted basis, or can be constructed from readily available plasmids in accordance with published procedures.

20 A "primer" is a nucleic acid fragment which functions as an initiating substrate for enzymatic or synthetic elongation of, for example, a nucleic acid compound.

The term "promoter" refers to a nucleic acid sequence that directs transcription, for example, of DNA to RNA. An  
25 inducible promoter is one that is regulatable by environmental signals, such as carbon source, heat, or metal ions, for example. A constitutive promoter generally operates at a constant level and is not regulatable.

"Recombinant DNA cloning vector" as used herein refers  
30 to any autonomously replicating agent, including, but not limited to, plasmids and phages, comprising a DNA molecule

-16-

to which one or more additional DNA segments can or have been incorporated.

The term "recombinant DNA expression vector" or "expression vector" as used herein refers to any recombinant DNA cloning vector, for example a plasmid or phage, in which a promoter and other regulatory elements are present thereby enabling transcription of an inserted DNA, which may encode a polypeptide.

The term "selectively binding" refers to the ability of FLINT polypeptides to bind FasL but not  $\text{TNF}\alpha$ .

"Substantially pure" used in reference to a peptide or protein means that said peptide or protein is separated from other cellular and non-cellular molecules, including other protein molecules. A substantially pure preparation would be about at least 85% pure; preferably about at least 95% pure. A "substantially pure" protein as described herein could be prepared by a variety of techniques well known to the skilled artisan, including, for example, the IMAC protein purification method.

The term "treatment" or "treating" as used herein, describes the management and care of a patient for the purpose of combating a disease, condition, or disorder and includes the administration of FLINT to prevent the onset of the symptoms or complications, alleviating the symptoms or complications thereof.

The term "vector" as used herein refers to a nucleic acid compound used for introducing exogenous or endogenous DNA into host cells. A vector comprises a nucleotide sequence which may encode one or more protein molecules. Plasmids, cosmids, viruses, and bacteriophages, in the

-17-

natural state or which have undergone recombinant engineering, are examples of commonly used vectors.

The various restriction enzymes disclosed and described herein are commercially available and the manner of use of said enzymes including reaction conditions, cofactors, and other requirements for activity are well known to one of ordinary skill in the art. Reaction conditions for particular enzymes were carried out according to the manufacturer's recommendation.

10

## II. FLINT binds Fas L and LIGHT

FLINT is a newly identified member of the TNFR superfamily. This family of receptors mediates a variety of biological effects of TNF ligands, including but not limited to cell proliferation, cell differentiation, immune regulation, inflammatory response, cytotoxicity, and apoptosis.

The FLINT polypeptide of the present invention is a soluble receptor comprising extracellular domains. FLINT polypeptide does not include any transmembrane domains and is, therefore, soluble. FLINT has also been called OPG3 (osteoprotegrin 3) or TNFRsol. FLINT is believed to be closely related to TNFR 6 $\alpha$  and TNFR 6 $\beta$  discussed in WO98/30694 claiming priority to U.S.S.N. 60/035,496 and TR4, discussed in EP 0861850A1.

Data presented below demonstrate that mFLINT binds to FasL and to LIGHT *in vitro*. FasL, as noted, is involved in triggering both an inflammatory response and in inducing apoptosis. The biological activity for LIGHT includes, but is not limited to, cell proliferation. LIGHT is a 29kDa

30

-18-

type II transmembrane TNF superfamily member protein produced by activated T cells. Mauri D.M., *Immunity*, 8:21 (1998). As with FasL, evidence links LIGHT with neutrophil infiltration and with apoptosis. Zhai et al., *J. Clin. Invest.* 102:1142-51, 1998. Thus, mFLINT-mediated inhibition of LIGHT activity is expected to be therapeutically useful substantially as described below for mFLINT-mediated inhibition of FasL, especially in immune modulation and cancer therapy.

10

### III. FLINT - Therapeutic Applications

The inventors have discovered that mFLINT prevents *in vitro* FasL-induced apoptosis of Jurkat cells. Anti-CD3 antibodies activate these cells and cause them to express FasL and undergo apoptosis. mFLINT also was effective in inhibiting *in vitro* anti-CD3-induced apoptosis of Jurkat cells, in a dose-dependent manner.

The present invention is applicable to the treatment and/or inhibition of disorders that may be associated with an inappropriate or abnormal interaction of FasL with Fas, because mFLINT prevents this interaction. This inappropriate interaction may arise, for example, due to increased expression or availability of FasL and/or Fas. Since it is known that FasL-Fas interaction induces apoptosis, the inventive methods further generally apply to disorders characterized by inappropriate or abnormal apoptosis.

In another embodiment the present invention relates to a method of preventing or treating conditions caused or exacerbated by FasL-Fas binding including FasL-mediated

30

-19-

apoptosis and/or a proinflammatory response, more particularly, a proinflammatory response caused by FasL induced neutrophil activation.

For instance, mFLINT may be employed in treating Down's syndrome. Enhanced apoptosis of neuronal cells may be implicated in Down Syndrome. In this regard, Seidi, et al., *Neuroscience Lett.* 260:9 (1999), reported elevated levels of Fas protein in the temporal lobe and cerebellum of adult patients with Down syndrome, suggesting that Fas-associated apoptosis may be an important feature of neurodegeneration in Down syndrome. It is expected, therefore, that treatment of a Down syndrome patient with mFLINT may be effective. In particular, the present inventors have shown that mFLINT binds FasL, prevents Fas-FasL binding, and inhibits apoptosis. Similarly, apoptosis has been linked to Alzheimer's disease and other neurodegenerative diseases. Administration of mFLINT is expected to decrease the enhanced apoptosis associated with Down syndrome, Alzheimer's disease and other neurodegenerative disorders.

The inventors have tested mFLINT in a variety of model systems that are indicative of various disease and injury states. Some exemplary disorders include, but are not limited to, antibody-dependent cytotoxicity, infection with human immunodeficiency virus, hemolytic uremic syndrome, allergies and bronchopulmonary dysplasia. Thus, the following section discusses diseases and injury states in the context of corresponding model systems.

The present invention also encompasses the production of transgenic animals that contain a FLINT transgene. In particular, the inventors have produced transgenic mice that produce measurable levels of mFLINT. Animals such as these

-20-

are useful for assessing the effects of mFLINT on a variety of disorders, detailed below, such as endotoxin-induced shock, cerebral ischemia, cardiac reperfusion injury, and damage induced by cancer therapies, such as therapeutic  
5 irradiation and chemotherapy.

**A. Treatment of Liver damage and inflammation with mFLINT**

FasL has been implicated in acute liver failure including but not limited to the damage caused to the liver in hepatitis. In a mouse model of acute hepatitis,  
10 administration of anti-FasL antibody to mice caused liver failure induced by apoptosis in hepatocytes and the animals die within hours. Kondo et al, 1997 Nature Medicine 3(4):409-413. See also Galle et al, J. Exp. Med, November 1995, 182:1223-1230.

15 Tsuji et al. (1998), Infect. Immun. 65:1892-1898, describes a model system for bacterially induced fulminant hepatitis. As detailed below, this system is predictive of a variety of disorders of the liver and other tissues. In the method of Tsuji et al., mice were injected with  
20 *Propionibacterium acnes*, and are subsequently injected with lipopolysaccharide (LPS). In normal mice, the first injection causes granuloma formation in the liver, while the second injection induces massive apoptosis and consequent liver damage. Tsuji et al. employed this method to  
25 implicate TNFRp55 in granuloma formation, and TNFRp55 and Fas in apoptosis.

Using a modified version of the animal model of Tsuji et al., the applicants have found that administration of mFLINT dramatically improves the survival rate of test  
30 animals. As indicated, many of the disorders treatable and/or preventable according to the present invention share

-21-

a FasL-Fas-related etiology. Thus, target diseases generally involve either a tissue inappropriately (e.g., temporally, qualitatively or quantitatively) expressing Fas or inappropriately coming into contact with FasL while  
5 expressing Fas. Many of these diseases follow a general pathological model of inappropriate upregulation of Fas, followed by FasL-mediated induction of apoptosis. The inappropriate expression of Fas (or FasL) may be induced, for example, by inflammatory insult, with certain  
10 inflammation-associated cytokines likely inducing expression.

For instance, it is known that the first-phase mononuclear infiltration results in the secretion of numerous cytokines, which may result in inappropriate Fas and/or FasL expression. It is known, for example, that IL-  
15 1, IL-1 and TNF- can induce Fas expression. This increased Fas expression may, in effect, sensitize the affected tissue to FasL-bearing effector cells, like natural killer cells and cytotoxic T cells. Contact of the FasL-bearing effectors with Fas-bearing targets induces apoptosis  
20 of the latter. In other words this hepatic injury model is an *in vivo* surrogate for certain disease characterized by (a) inflammatory insult and/or (b) FasL-Fas-mediated apoptosis and/or necrosis.

Consistent with this model is the pathology of graft  
25 versus host disease (GVHD) which is very common, for example, in autologous bone marrow transplantation. GVHD results from the presence of host-reactive T cells that destroy host tissues, at least in part via FasL-mediated apoptosis. GVHD is typically divided into two phases,  
30 termed afferent and efferent. The afferent phase is characterized by recognition of host antigens and

-22-

proliferation of donor T cells. In efferent phase tissues like the skin, liver and gastrointestinal tract become inflamed, and it is characterized by mononuclear cell infiltration and histopathological damage. Experiments using FasL-defective mice, in fact, show that FasL plays a key role in the development of hepatic, cutaneous and lymphoid organ damage in GVHD. Hence, GVHD follows the paradigm of inflammatory insult and Fas-mediated apoptosis.

Hashimoto's thyroiditis (HT) also fits this paradigm. HT results from an autoimmune response against the thyroid follicular cells. Normal thyrocytes produce FasL and express negligible levels of Fas. In HT, however, the resulting inflammation results in IL-1 secretion by activated macrophages, which then induces the thyrocytes to produce Fas. This sets up a fatal FasL-Fas autocrine loop that results in apoptosis.

Similarly, oxidized low density lipoprotein (OxLDL), associated with atherosclerotic lesions, promotes a chronic inflammatory response. While the vascular endothelium normally expresses both FasL and Fas, absent insult by OxLDL it does not undergo apoptosis. Upon treatment with OxLDL, however, FasL expression is increased and the endothelial cells undergo apoptosis.

Chronic renal failure is correlated secretion of IL-1 and TNF- $\alpha$ , both of which have been shown to induce Fas expression in renal tubular epithelial cells. This disorder is characterized by the depletion of tubular epithelial cells, via the FasL-Fas apoptosis pathway. Moreover, the same cytokine-mediated induction of Fas expression in tubule cells is observed in the endotoxic shock (sepsis) model of acute renal failure.



-23-

Acute liver failure can be found in viral infections affecting the liver, bacterial infections affecting the liver, hepatitis, hepatocellular injury and/or other conditions where hepatocytes undergo massive apoptosis.

5 Galle et al, (J. Exp. Med., November 1995, 182:1223-1230), for example, found that FasL-Fas mediated cell death played a role in liver failure in humans. Galle et al. knew that FasL-Fas mediated apoptosis was a mechanism to eliminate senescent hepatocytes. For example, it was found in liver  
10 regeneration, during involution of liver after cessation of treatment with lipophyllic compounds (e.g., phenobarbitol) and during viral infection. Through experiments they found that during viral hepatitis, activated T cells attacked hepatocytes. Fas is constitutively expressed in  
15 hepatocytes. That is, FasL was expressed in T cells upon activation. It is thought that activated T cells might kill hepatitis B (HBV) antigen-expressing hepatocytes in an effort to clear HBV from the liver.

The inventors' success using a modified model of Tsuji  
20 further indicates that mFLINT is useful in treating and preventing sepsis. Sepsis is characterized by the presence of one or more pathogenic organisms, or their toxins, in the blood or tissues. Furthermore, sepsis is characterized by a systemic inflammatory response to infection that is  
25 associated with and mediated by the activation of a number of host defense mechanisms including the cytokine network, leukocytes, and the complement and coagulation/fibrinolysis systems. See Mesters, et al. *Blood*, 88:881 (1996).

It is known that endotoxin(s) induce tissue necrosis  
30 factor (TNF), which in turn induces both an inflammatory response and FasL, thereby promoting liver failure and

-24-

death. Attempts to use antibodies drawn to TNF- in treating sepsis, however, have uniformly failed. This is likely do to the fact that, aside from its effects on the pathology of sepsis, TNF- is a more global mediator of normal immune function. In addition, as demonstrated below in the examples, TNF inhibitors show little usefulness in treating post-insult sepsis; rather they are more useful prophylactically.

Data presented below in the examples confirm that mFLINT is more effective than TNF inhibitors in promoting survival in an animal sepsis model. In fact, these data show that, whereas TNF inhibitors are efficacious only if used in a pre-treatment protocol (*i.e.*, before LPS/galactosamine administration), mFLINT may be used in a post treatment regimen. This is clinically significant because the patient will virtually always present after endotoxic exposure. In other words, these data demonstrate that mFLINT is useful in both preventative and ameliorative regimens, in marked contrasted to TNF inhibitors. Minimally, these data indicate that mFLINT may be administered at a later time in disease progression than anti-TNF; it is thus expected to avoid the problems that were encountered with TNF inhibitors in the clinic.

Moreover, these data demonstrate that mFLINT has a therapeutic effect on at least two levels. Endotoxin is known to induce TNF, which directly induces both an inflammatory response and FasL. The data of Tsuji *et al*, *supra*, indicate that these represent two distinct pathways, both of which contribute to liver failure and death. In particular, their data using Fas-deficient mice strongly suggested that a substantial amount of the damage in their

-25-

model is done through a Fas-independent, TNF-dependent pathway. The data presented below, however, demonstrate that mFLINT, an inhibitor of the Fas-dependent pathway, can inhibit all damage induced in the model, Fas-dependent or  
5 not. Thus, even if they represent different pathways, mFLINT can inhibit the damage mediated by both the inflammatory response and the induction of FasL.

In addition, it is contemplated that mFLINT may be beneficially used in combination with other agents useful in  
10 treating sepsis. For instance, U.S. Patent No. 5,009,889 demonstrates that activated protein C (aPC) is effective in treating a baboon sepsis model. Such a model may be employed to determine appropriate combination treatment protocols using, for example, mFLINT and aPC. aPC may  
15 prevent the sepsis-associated disseminated intravascular coagulation and widespread deposition of fibrin in the microvasculature, which is an early manifestation of sepsis/septic shock.

Thus, based on the improved survival rate observed in  
20 mFLINT-treated animals, the inventors contemplate that one or more of the following diseases can be treated with mFLINT: acute respiratory distress syndrome (ARDS); goiter; hepatocellular injury (viral, bacterial, cancer, trauma); mononucleosis; mucocites (inflammation of the mucosa);  
25 pancreatitis; periodontal disease/gingivitis; renal failure; satellite organ failure; systemic inflammatory response syndrome (SIRS); surgery; vascular bleeds; vascular leak syndrome; whole organ transplant; multiple organ dysfunction (MODS); coronary artery bypass grafting; allograft  
30 rejection; transplant rejection; infection (e.g., microbial, pneumonia, tissue necrotic infection, viral); bone loss in

-26-

rheumatoid arthritis; Hashimoto's thyroiditis; viruses, including hepatitis C (HCV), hepatitis B (HBV), Ebola (hemolytic fever) and Epstein-Barr (EBV); cutaneous inflammation; psoriasis; inflammatory bowel disease; ulcerative colitis; Crohn's disease; atherosclerosis; end stage renal disease; and sepsis.

More particularly, the invention contemplates methods for treating disorders associated with inflammation. The skilled artisan will recognize that these disorders include but are not limited to, GVHD, Hashimoto's thyroiditis, ARDS, heptatocellular injury (viral, bacterial, cancer, trauma); mucocites (inflammation of the mucosa); pancreatitis, periodontal disease/gingivitis; renal failure; satellite organ failure; systemic inflammatory response syndrome (SIRS); whole organ transplant; multiple organ dysfunction (MODS); coronary artery bypass grafting; allograft rejection; transplant rejection; infection (e.g., microbial, pneumonia, tissue necrotic infection, viral); rheumatoid arthritis; viral infections, including hepatitis C (HCV), hepatitis B (HBV), Ebola (hemolytic fever) and Epstein-Barr (EBV); cutaneous inflammation; inflammatory bowel disease; atherosclerosis; and sepsis.

**B. Treatment of Ischemia and Reperfusion with mFLINT**

Ischemia is characterized by the reduced blood flow to a tissue, which results in the accumulation of a variety of toxic metabolites. These metabolites contribute to cell death resulting from necrosis and/or apoptosis. Perhaps surprisingly, when a tissue is reperfused, apoptotic damage is increased. This is likely due to the generation of free radicals and other toxins through the reaction of ischemia-induced metabolites with serum components. In any event,

-27-

studies have shown that the apoptotic damage results at least in part from Fas-mediated pathways. Experiments presented below demonstrate that mFLINT can be used to block ischemia/reperfusion injury, likely by inhibiting FasL.

5       The usefulness of mFLINT compositions in treating or preventing disorders associated with cerebral ischemia, like stroke and head trauma, is confirmed by data presented below. There, a model is presented that uses a live gerbil model of global stroke. Cerebral ischemia was induced by a  
10   transient occlusion of the common carotid arteries followed by a treatment of mFLINT or a vehicle control. Data show that the mFLINT-treated gerbils retained markedly more living neurons than the vehicle treated control group.

      These data are consistent with studies comparing the  
15   extent of infarct brain tissues in mice lacking functional Fas receptors after a cerebral ischemia insult, which found that significantly less damage than normal controls. Rosenbaum, et al., *Annals of Neurology*, 44(3), 441, 1998. It is, therefore, anticipated that mFLINT will interfere the  
20   normal signaling process of the Fas receptor in a patient that has undergone a cerebral ischemia episode thus protecting the brain tissues from deterioration.

      Moreover, Fas-mediated apoptosis has been linked to ischemia reperfusion injury in a myocardial infarction  
25   model. Kajstura et al., *Laboratory investigation* 74:86-107 (1996). Thus, the foregoing gerbil model likely is predictive of generalized ischemia reperfusion injury. In support of this, the inventors conducted in an *in vitro* assay using cardiac myocytes. In particular, cardiomyocytes  
30   were incubated under hypoxia conditions for 8 hours, followed by 16 hours of culture under normal O<sub>2</sub>, which

-28-

mimics the hypoxic conditions found in ischemic cardiac tissue. Experimental results showed that treating the cardiomyocytes with mFLINT protected the cells against apoptosis induced by hypoxia. In particular, treatment with  
5 10 µg/ml mFLINT resulted in a 90% inhibition of hypoxia-induced apoptosis. FasL-induced apoptosis of cardiomyocytes also was inhibited by treatment with mFLINT.

Based on the foregoing observations and data, the inventors contemplate that one or more of the following  
10 diseases can be treated with mFLINT: stroke; spinal cord ischemia; eclampsia/preeclampsia; reperfusion injury; myocardial infarction, its acute, subacute and chronic sequelae and related clinical syndromes, including but not limited to congestive heart failure. Thus, FLINT is useful  
15 in promoting myocardial salvage and preventing decompensatory myocardial hypertrophy.

Reperfusion injury may be in a number of tissues and result from a variety of insults, including the bowel, burns, cardiac bypass machine injury, hepatic (trauma-  
20 induced), hemolytic fever (Ebola), infant toxicity (hyperoxia incubators with high O<sub>2</sub> content), limb crush lung injury/ARDS, organ transplant, multiple trauma (e.g. from car accidents), protection of vascular beds, vascular organ failure usually present in sepsis and other diseases. These  
25 observations further indicate that mFLINT is useful in the prevention/attenuation of apoptosis of cardiac myocytes following acute myocardial infarction, and generally in preventing damage to cardiac and other tissue due to hypoxia.

30 In the case of organ preservation in preparation for harvesting, for instance, mFLINT is useful prophylactically

-29-

to prevent the apoptosis associated with ischemia  
reperfusion injury to the organ once it is removed from the  
donor. In one embodiment of this aspect of the invention,  
FLINT is administered to the organ donor prior to harvesting  
5 the organ. After harvesting the organ, FLINT is added to a  
suitable medium for transport/storage of the organ.  
Alternatively, the harvested organ is perfused with a medium  
containing FLINT prior to transplantation into a recipient.  
Suitable media for this purpose are known, for example, the  
10 media disclosed in EP 0356367 A2, herein incorporated by  
reference. The method may also include treating the  
transplant recipient with FLINT prior to and/or after the  
transplant surgery.

A typical method involves pre-treating the organ donor  
15 with an effective amount of mFLINT prior to organ  
harvesting. Alternatively, or conjunctively, the harvested  
organ may be perfused or bathed in a mFLINT-containing  
solution. This method may be employed, for example, with  
kidney, heart, lung and other organs and tissues.

20 In another example, mFLINT is useful in treating the  
ischemia reperfusion injury associated with thrombus  
formation or hypercoagulation. Thus, mFLINT may be  
administered in conjunction with thrombolytic agents (e.g.,  
urokinase and streptokinase) and/or antithrombotics, like  
25 activated protein C (aPC). U.S. Patent No. 5,350,578  
describes the antithrombotic activity of aPC. The baboon  
animal model disclosed therein may be used to ascertain a  
beneficial dosing regimen for the contemplated combination  
therapy. In this regard, it is expected that the following  
30 disorders may be treated with mFLINT, with or without the  
addition of aPC: eclampsia and preeclampsia, which are

-30-

characterized by a state of increased coagulopathy; HELLP (preeclampsia complicated by thrombocytopenia, hemolysis, and disturbed liver function); HITS (heparin-induced thrombocytopenia); disseminated intravascular coagulation (DIC); burns; and thromboembolic complications that result from surgery.

As noted above, studies show that a great deal of apoptotic injury is induced upon reperfusion. This injury appears to be due at least in part to oxidative damage, since antioxidants can diminish the injury. Likewise, it has been found that Vitamin E, an antioxidant, can be used as a meat preservative, perhaps acting to prevent the same type of injury, which causes spoilage. Administering Vitamin E to pigs several days before slaughter has been shown to increase shelf-life of pork. Hence, it is contemplated that mFLINT is similarly useful to preserve tissues and organs. This would be particularly useful for increasing shelf-life of organs and meats for human consumption.

20

#### C. Treatment of Hematopoietic Disorders with mFLINT

As discussed in this application, mFLINT inhibits Fas/FasL interaction, thereby preventing apoptosis. As discussed below, the Fas/FasL interaction has been implicated in the process of hematopoiesis. Therefore, the present application contemplates mFLINT-based therapeutic methods for improving hematological recovery from a variety of treatments and disorders, including bone marrow transplantation, chemotherapy, radiotherapy, aplastic anemia, and myelodysplastic syndrome, pancytopenic conditions. The present invention also covers the mFLINT-

30



-31-

based treatment of any clinical condition which requires expansion of bone marrow cell lineages, alone or in combination with other hematological growth factors. Such growth factors include, but are not limited to, erythropoietin (EPO), FLT-3 ligand, thrombopoietin (TPO), stem cell factor (SCF), granulocyte colony stimulating factor (G-CSF), and granulocyte-macrophage colony-stimulating factor (GM-CSF).

Autologous and heterologous bone marrow transplantation are commonly used to treat a multitude of neoplastic diseases. In autologous transplantation, bone marrow cells are removed from the patient to be treated, and are cultured *in vitro*. Following chemotherapy and/or radiation therapy, the cells are re-introduced to the patient. In heterologous transplantation, the bone marrow cells are donated by a second individual. The chemotherapy or radiotherapy that is used to treat such diseases results in a dramatic myelosuppression from damage to the bone marrow compartments and intestinal epithelium, leaving the patient immunosuppressed and susceptible to invasion by microorganisms. The treatment of transplanted bone marrow cells with hematopoietic cytokines has been shown to result in improved recovery of the myeloid and erythroid lineages of the blood, although there typically is toxicity associated with the use of these cytokines and one cytokine typically does not enhance recovery of all lineages. Moore, M.A.S. *Blood* 78:1 (1991); Metcalf, D. *Science* 254: 529 (1991).

Radiation or chemotherapy has been shown to induce apoptosis of bone marrow cells and intestinal epithelial cells. The Fas/FasL pathway is known to induce apoptosis in

-32-

susceptible cells. Recent reports describe the involvement of Fas/FasL in hematopoiesis. For a review, see Niho, et al.; *Current Opinion in Hematology*, 5: 163 (1998). Fas is expressed on CD34+ progenitor cells and these cells are susceptible to the apoptotic effects of Fas stimulation. Nagafuji et al., *Blood*, 86: 883-889 (1995), Sato et al., *Br J Haematol*, 97: 356 (1997). Yamane et al., *Eur J Haematol*, 58:289 (1997). Moreover, *in vitro* expansion of hematopoietic progenitor cells with cytokines has been shown to upregulate functional Fas expression and it is currently thought that Fas may play an *in vivo* role in hematopoietic homeostasis as a negative regulator. Takenaka et al., *Blood*, 88: 2871 (1996), Stahnke et al., *Exp. Hematol.* 26: 844 (1998).

15

#### 1. mFLINT and Radiation and Chemotherapy

The inventors have shown that mFLINT improves the recovery of bone marrow progenitor cells that have been exposed to irradiation. In particular, mice were irradiated, and bone marrow cells were removed and cultured *in vitro* with the cytokines IL-6 and CSF. The addition of mFLINT to the culture medium dramatically increased the recovery of bone marrow progenitor cells, namely progenitors of erythroid cells, granulocyte-macrophage cells, and erythroid monocyte megakaryocyte cells. The inventors also have shown that mFLINT improves the survival of bone marrow cells that were harvested from mice treated with 5-fluorouracil (5-FU). Anti-FasL antibody also increased the survival of the cells taken from 5-FU-treated mice.

The present invention thus encompasses the use of mFLINT as a radioprotective and/or chemoprotective agent.

-33-

mFLINT is useful for enhancing the *in vitro* expansion and maturation of bone marrow progenitor cells prior to autologous or heterologous transplantation. mFLINT also is useful for promoting the expansion and maturation of  
5 progenitor cells, after a patient has been treated with radiation or chemotherapy. In this regard, administration of mFLINT to a patient is expected to promote the expansion and maturation of progenitor cells when that patient has been treated with a cancer therapy (chemo- or radio-  
10 therapy). mFLINT also is expected to promote the expansion and maturation of progenitor cells in patients treated with cancer therapy, and myelosuppressive agent, which prevents cycling of progenitor cells during cancer therapy.

As noted above, cytokines are used, *in vitro* and *in vivo*, to enhance the expansion of erythroid and myeloid cell  
15 lineages in patients receiving cancer therapies. The present inventors have shown that mFLINT improves the expansion of cells that are treated with cytokines. Accordingly, the present invention contemplates the use of mFLINT, in combination with one or more cytokines, to expand  
20 bone marrow progenitor cells, *in vitro* and *in vivo*. Granulocyte-macrophage colony stimulating factor (GM-CSF) is used to improve recovery from myelosuppression that results from chemo- and radio-therapy. Accordingly, the present  
25 invention contemplates the use of a combination of mFLINT and GM-CSF to improve recovery from myelosuppression.

The inventors also have shown that anti-FasL antibody also improved the recovery of bone marrow cells from treatment with 5-FU. Accordingly, the present invention  
30 contemplates the use of anti-FasL to improve recovery of bone marrow cells, following chemo- or radio-therapy. Such

-34-

an antibody can be used *in vitro* to expand bone marrow cells for heterologous or autologous transplantation. An anti-FasL antibody also can be administered *in vivo* to improve recovery from myelosuppression that results from  
5 chemotherapy, radiation therapy, and/or administration of myelosuppressive agents.

Systemic administration would be appropriate when treating patients with mFLINT. As used in this application, administration of mFLINT "with" chemotherapy or irradiation  
10 encompasses all of the following modes of administration of mFLINT: (1) pretreatment with mFLINT, followed by irradiation or chemotherapy; (2) simultaneous administration of mFLINT and chemo- or radiation therapy; (3) chemotherapy or irradiation first, followed by mFLINT administration; (4)  
15 mFLINT pretreatment, followed by chemo- or radiation therapy, followed by administration with mFLINT. The invention encompasses the use of mFLINT in one or more of the foregoing modes of treatment.

## 2. Treatment of peripheral cytopenias

20 The present invention also contemplates the treatment of peripheral cytopenias that are associated with hematopoietic disorders, such as aplastic anemia and myelodysplastic syndrome. Fas/FasL-mediated apoptosis has been shown to suppress lymphopoiesis. Yasutomo, et al. *J. Immunol.* 157: 1981 (1996). Increased Fas expression is  
25 observed in progenitor cells from patients with aplastic anemia. Young, N.S., *Eur. J. Hematol.* 60 (supp): 55 (1996). Increased Fas expression also has been observed in progenitor cells from patients suffering from  
30 myelodysplastic syndrome.

-35-

Furthermore, Maria, et al. reported that Fas is rapidly upregulated in early erythroblasts and is expressed at high levels thorough terminal maturation. Blood 93:796 (1999). Maria, et al. also reported that immature blood cells are  
5 EPO-dependent and express a functional Fas molecule. In the presence of FasL-producing mature erythroblasts, the immature cells undergo apoptosis, unless exposed to high levels of EPO. Maria, et al. suggest that Fas/FasL system, in combination with EPO, contribute to erythropoietic  
10 homeostasis.

While not intending the bound to any particular theory, the reduction in hematopoiesis and resultant cytopenias may be a direct result of apoptosis, which may in turn be mediated through upregulation of Fas and Fas ligand. As  
15 described elsewhere in this application, mFLINT inhibits Fas/FasL binding, inhibiting apoptosis, and mFLINT also enhanced the growth and maturation of hematopoietic progenitor cells. Therefore, the use of mFLINT is expected to suppress apoptosis suffering from hematopoietic  
20 disorders, leading to amelioration of one or more of the symptoms associated with these disorders. Moreover, because immature hematopoietic cells are susceptible to FasL-induced apoptosis, and because individuals suffering from a lack of differentiated hematopoietic cells, the administration of  
25 mFLINT to such individuals is expected to ameliorate these maturation defects, by inhibiting FasL-induced apoptosis of immature hematopoietic cells.

Therapy of individuals suffering from hematopoietic disorders can be carried out by direct administration of  
30 mFLINT to the affected individual, or by the use of mFLINT

-36-

in vitro to treat bone marrow cells that are then transplanted into the diseased individual.

Therapy with mFLINT can be augmented by administration of EPO and other compounds that promote the growth and/or maturation of hematopoietic progenitor cells. Cytokines are known to stimulate the growth of hematopoietic progenitor cells. Accordingly, the invention also encompasses the use of a combination of mFLINT and one or more cytokines to promote growth and differentiation of such progenitor cells in individuals suffering from diseases such as thrombocytopenia and myelodysplastic syndrome.

### 3. Gene Therapy

FLINT also can be used in conjunction with gene therapy. When hematopoietic progenitor cells are to be transplanted into an individual, they are transfected with a suitable transgene, and also are treated with mFLINT, optionally in combination with one or more cytokines. Following culturing of the cells, they are transplanted into an individual. Such gene therapy will be useful in imparting desirable properties to the blood cells.

### D. Protection of Innocent Bystander Tissues with mFLINT

It is well known that many cell-damaging therapies, such as chemotherapy and therapeutic irradiation used to treat cancer, cause damage and apoptosis of so-called "innocent bystander" tissues. As used in this application, an "innocent bystander" tissue is a non-diseased tissue that is damaged by pharmacological, radiation or device-assisted therapies. These tissues include, but are not limited to, the gastric epithelium (including the epithelium lining the oral cavity, esophagus, stomach and intestinal tract), the

-37-

lung epithelium, blood cells (lymphocytes, monocytes, T cells, B cells, bone marrow cells, hematopoietic progenitor cells, neutrophils, eosinophils, mast cells, platelets), renal epithelium, and hair follicles. Many of these tissues  
5 are characterized by rapid growth and/or turnover of their constituent cells.

"Cell damaging" therapies include, but are not limited to chemotherapy (e.g. cisplatin, doxorubicin, mitomycin C, camptothecin, and fluorouracil and other nucleoside  
10 analogs), therapeutic irradiation, laser treatment, and administration of inhaled toxins (such as bleomycin). Physical manipulations that are associated with device-assisted therapies, such as the physical removal of a blockage from a coronary artery, also can cause damage to  
15 innocent bystander tissue. While these cell-damaging therapies are useful because they damage and kill diseased tissue, as noted above, they have the undesirable side effect of damaging and inducing apoptosis of "innocent bystander" tissue.

20 As discussed elsewhere in this application, mFLINT blocks Fas/FasL interactions. It also has been demonstrated herein that mFLINT inhibits FasL-induced apoptosis *in vitro*, and that mFLINT improved the survival of bone marrow cells following therapeutic irradiation and chemotherapy.  
25 Therefore, it is an object of the present invention to ameliorate damage to innocent bystander tissues that is caused by cell-damaging therapy.

Cancer therapies such as irradiation and chemotherapy have been shown to induce apoptosis in intestinal epithelial  
30 cells. It is known that many chemotherapeutic agents work by inducing apoptosis. See, e.g., Micheau, et al. *J. Natl.*

-38-

Can. Inst. 89:783 (1997). This apoptosis, combined with the myelosuppression that is induced by irradiation and chemotherapy, permits massive opportunistic infection.

There currently are no effective therapies for managing the  
5 disruption of intestinal function that is associated with chemotherapy and therapeutic irradiation. As noted above, treatment with cytokines has been shown to improve recovery from myelosuppression, but cytokines can produce undesirable toxicity.

10 Therefore, it is expected that mFLINT will ameliorate damage to the intestinal epithelium that is induced by irradiation and/or chemotherapy. While not desiring to be bound to any particular theory, it is expected that mFLINT will ameliorate this damage by inhibition of apoptosis  
15 and/or inflammation of the intestinal epithelium.

Fas/FasL interactions have been implicated in chronic gastritis. It has been shown that Fas and FasL expression are increased in gastric epithelial cells from individuals suffering from chronic gastritis. See Rudi, et al., *J.*  
20 *Clin. Invest.* 102:1506 (1998). Rudi also reported that *Helicobacter*-infected gastric epithelial cells have increased levels of FasL (CD95 ligand) and Fas (CD95 receptor), and that *Helicobacter*-induced apoptosis was reduced by blocking FasL with anti-APO-1 antibody. Thus, it  
25 is expected that mFLINT may be effective in inhibiting (i) gastritis that is induced by *Helicobacter* and, as mentioned above, (ii) cell-damaging therapies, such as chemotherapy and irradiation.

Furthermore, patients receiving high-dose chemotherapy  
30 (HD-CT) are at risk of severe mucositis, which is characterized by apoptosis and inflammation of the gastric



epithelium. Wymenga, et al., *Br. J. Cancer* 76:1062 (1997) studied mucositis of in the oral cavity of breast cancer patients treated with HD-CT. The percentage of viable epithelial cells appearing in an oral wash increased in  
5 patients treated with HD-CT, suggested a desquamation of the upper oral mucosal layer. Also, there was a higher percentage of immature cells in the oral mucosa of the HD-CT patients.

Other innocent bystander tissues also may benefit from  
10 mFLINT therapy. For example, Fas/FasL-induced apoptosis has been implicated in bleomycin-induced apoptosis and fibrosis in lung epithelium. Hagimoto, et al. *Am. J. Respir. Cell. Mol. Biol.* 16:91 (1997). That study showed that in alveolar epithelial cells treated with bleomycin, Fas mRNA was  
15 upregulated, and FasL mRNA was upregulated in infiltrating lymphocytes. Therefore, the applicants expect that mFLINT therapy may ameliorate damage in lung tissue, such as epithelium, that is induced by cell-damaging therapies.

20 **E. Treatment of Cancer with mFLINT**

The present inventors have found that administration of mFLINT to mice causes a reduction in the volume of tumors that are produced when mice are injected with melanoma cells. This effect of mFLINT may be mediated through  
25 mFLINT/FasL binding and the consequent inhibition of FasL-mediated T-cell apoptosis, permitting a T-cell mediated immune response to facilitate destruction of the tumor. In this regard, it is known that FasL is expressed in melanomas, human colorectal cancers, hepatocellular cancers,  
30 astrocytomas and lung cancer. O'Connell, et al. *Immunology Today* insert\*, Chappell, et al. *Cancer Immunol. Immunother.*

-40-

47:65 (1998). It also has been reported that colon cancer cell lines express FasL and can kill T cells in vitro by inducing FasL-mediated apoptosis. O'Connell. Therefore, the melanoma tumors observed in the present application may  
5 express FasL, triggering FasL-mediated death of infiltrating T-cells that express Fas. However, other mechanisms may also be responsible for the tumor reduction that has been demonstrated in the present application.

Based on the success achieved by the present inventors  
10 with mFLINT-mediated reduction of tumor volume, it is expected that other types of cancer also can be successfully treated with mFLINT. These cancers include, but are not limited to, astrocytoma, colon cancer, esophageal cancer, lung cancer, melanoma and hepatocellular cancer. Other  
15 cancers that can be treated with mFLINT include bladder cancer and ovarian cancer.

#### **F. Treatment of Autoimmune Disease with mFLINT**

Both Type I (insulin-dependent) and Type II (non-  
20 insulin dependent) diabetes are characterized by hyperglycemia. As used herein, the term "hyperglycemia," which is well known in the art, describes a condition characterized by a blood glucose level that is higher than that found in a normal human. Normal human fasting blood  
25 glucose levels are less than 110 mg/dL. The Expert Committee on the Diagnosis and Classification of Diabetes Mellitus, *Diabetes Care*, 21 (Suppl. 1), S5-S19 (1992).

Type I insulin-dependent diabetes (IDDM) is a chronic autoimmune disorder involving destruction of pancreatic  
30 (insulin-producing)  $\beta$  cells via the FasL-Fas pathway. It is likely that this destructive pathway is activated by the

-41-

inflammatory process. Normal  $\beta$  cells do not express Fas, but Fas is expressed in these cells during insulinitis. It is thus likely that lymphocyte-associated FasL contributes to the local destruction of these Fas<sup>+</sup> cells via a FasL-Fas interaction. To the extent that mFLINT is useful in antagonizing this pathway, it is useful in preventing or treating IDDM.

Multiple sclerosis (MS) is a degenerative inflammatory demyelinating disease. Fas has been found among the oligodendrocytes located along the lesion margin and in the adjacent white matter in both acute and chronic MS, whereas normal tissue shows little or none. It is likely that this abnormal expression of Fas is induced by cytokines released during the inflammatory process. Moreover, FasL-expressing cells have been co-localized with Fas-expressing apoptotic oligodendrocytes, suggesting involvement of this pathway in the pathology of this disease. Lupus, another autoimmune disease, also can be treated with mFLINT.

#### **G. Treatment of Osteoporosis with mFLINT**

Data presented below in the examples indicate that mFLINT is useful in preventing or treating disorders of bone loss, like osteoporosis. mFLINT also can be used to inhibit bone resorption. These data are consistent with data showing the naturally-occurring secreted member of the TNFR superfamily was recently reported as having a role in regulating bone resorption. Simonet et al. *Cell* 89:309(1997) (termed osteoprotegrin (OPG); Tsuda et al. *Biochem. and Biophys. Res. Comm.* 234:137 (1997) (termed osteoclastogenesis inhibitory factor (OCIF))), functioning essentially as inhibitors of differentiation of bone-resorbing osteoclasts. Consistently, studies have directly linked Fas to osteoblast

-42-

apoptosis. Jilka et al., 1998, J. Bone & Mineral Res.  
13:793-802; Kawakami et al., 1997, J. Bone & Mineral Res.  
12:1637-46.

5 Acute Lung Injury and Acute Respiratory Distress Syndrome

Acute lung injury (ALI) and acute respiratory distress  
syndrome (ARDS) represent disease entities that differ only  
in the severity of the hypoxemia present at diagnosis. A  
widely accepted parameter for diagnosis is the PaO<sub>2</sub> to FiO<sub>2</sub>  
10 ratio, according to which ARDS patients manifest a ratio of  
less than or equal to 200 mm Hg, whereas ALI patients  
exhibit values of less than or equal to 300 mm Hg. ARDS  
represents a more severe form of ALI. Numerous mediators are  
likely to contribute to the pathogenesis of ARDS/ALI with  
15 neutrophils playing a prominent role. While multiple  
precipitating factors are probable in the development of  
ARDS, both direct and indirect, the major cause appears to  
be sepsis and the systemic inflammatory response syndrome,  
accounting for approximately 40% of cases. Mortality in ARDS  
20 is high approximating 40%, with most deaths occurring within  
the first 2 to 3 weeks. There is no currently available,  
approved pharmacologic therapy for ARDS and treatment at  
present is limited to aggressive supportive care.

There is evidence that ARDS may be mediated by soluble  
25 FasL/Fas interaction in humans (Matute-Bello et al., J.  
Immunol. 163, 2217-2225, 1999). FLINT, by binding to FasL,  
could inhibit FasL-mediated apoptosis of pneumocytes and/or  
endothelial cells, thus inhibiting or preventing the  
progression from acute inflammatory insult to ALI, and from  
30 ALI to ARDS.

-43-

In one embodiment the present invention relates to the use of FLINT to treat ALI and/or ARDS comprising the administration of a therapeutically effective amount of FLINT to a person in need thereof.

5

#### IV. THERAPEUTIC FORMULATIONS OF mFLINT

The mFLINT polypeptide composition will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with mFLINT polypeptide alone), the site of delivery of the mFLINT polypeptide composition, the method of administration, the scheduling of administration, and other factors known to practitioners.

An effective amount of polypeptide results in a statistically significant modulation of the biological activity of the selected TNFR family ligand, for example, FasL or LIGHT. The biological activity for FasL includes, but is not limited to, apoptosis. The biological activity for LIGHT includes, but is not limited to, cell proliferation. LIGHT is a 29 kDa type II transmembrane TNF superfamily member protein produced by activated T cells. Mauri d.M., Immunity, 8:21-30, January 1998.

Further, an effective amount may also be determined by prevention or amelioration of adverse conditions or symptoms of diseases, injuries or disorders being treated. The "therapeutically effective amount" of mFLINT polypeptide for purposes herein is thus determined by such considerations. It should be noted that mFLINT is an immunomodulator and that a common observation with such substances is a bell-

-44-

shaped dose-response curve. Such a phenomenon is well known in the art and it is within the skill of the clinician to take this into account in adjusting the therapeutically effective amount of mFLINT accordingly.

- 5 As a general proposition, the total pharmaceutically effective amount of mFLINT polypeptide administered parenterally per dose will be in the range of about 1  $\mu\text{g/kg/day}$  to 10  $\text{mg/kg/day}$  of patient body weight, more particularly 2-8 $\text{mg/kg}$ , preferably 2-4 $\text{mg/kg}$ , most preferred 10 2.2 $\text{mg/kg}$  to 3.3  $\text{mg/kg}$  and finally 2.5  $\text{mg/kg}$ . However, as noted above, this will be subject to therapeutic discretion. Preferably, this dose is at least 0.01  $\text{mg/kg/day}$ .

- If given continuously, the mFLINT polypeptide is typically administered at a dose rate of about 0.1 15  $\mu\text{g/kg/hour}$  to about 50  $\mu\text{g/kg/hour}$ , either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses 20 to occur appears to vary depending on the desired effect.

- Pharmaceutical compositions containing the mFLINT of the invention may be administered using a variety of modes that include, but are not limited to, oral, rectal, intracranial, parenteral, intracisternal, intravaginal, 25 intraperitoneal, topical, transdermal (as by powders, ointments, drops or transdermal patch), buccally, or as an oral or nasal spray. By "pharmaceutically acceptable carrier" is meant a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation 30 auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include but are not

-45-

limited to, intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion. Implants comprising mFLINT also can be used.

The mFLINT polypeptide is also suitably administered by  
5 sustained-release systems. Suitable examples of sustained-release compositions include semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules. Sustained-release matrices include polylactides (U.S. Pat. No. 3,773,919, EP 58,481),  
10 copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman, U. et al., *Biopolymers* 22:547-556 (1983)), poly (2-hydroxyethyl methacrylate) (R.Langer et al., *J. Biomed. Mater. Res.* 15:167-277 (1981), and R. Langer, *Chem. Tech.* 12:98-105 (1982)), ethylene vinyl acetate (R. Langer et al.,  
15 Id.) or poly-D- (-)-3-hydroxybutyric acid (EP 133,988). Sustained-release mFLINT polypeptide compositions also include liposomally entrapped mFLINT polypeptides. Liposomes containing mFLINT polypeptides are prepared by methods known per se: DE 3,218,121; Epstein et al., *Proc. Natl. Acad. Sci.*  
20 (USA) 82:3688-3692 (1985); Hwang et al., *Proc. Natl. Acad. Sci.* (USA) 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EDP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S. Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about  
25 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the optimal TNFR polypeptide therapy.

For parenteral administration, in one embodiment, the  
30 mFLINT polypeptide is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable

-46-

form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation. For example, the formulation preferably does not include oxidizing agents and other compounds that are known to be deleterious to polypeptides.

Generally, the formulations are prepared by contacting the mFLINT polypeptide uniformly and intimately with liquid carriers or finely divided solid carriers or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten residues) polypeptides, e.g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose



-47-

or its derivatives, glucose, manose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic surfactants such as polysorbates, poloxamers, or PEG.

5       The mFLINT polypeptide is typically formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml, preferably 1-10 mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the foregoing excipients, carriers, or stabilizers will result in the  
10   formation of mFLINT polypeptide salts.

FLINT polypeptides to be used for therapeutic administration must be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e.g., 0.2 micron membranes). Therapeutic mFLINT  
15   polypeptide compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

FLINT polypeptides ordinarily will be stored in unit or  
20   multi-dose containers, for example, sealed ampoules or vials, as an aqueous solution or as a lyophilized formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1% (w/v) aqueous mFLINT polypeptide  
25   solution, and the resulting mixture is lyophilized. The infusion solution is prepared by reconstituting the lyophilized mFLINT polypeptide using bacteriostatic Water-for-Injection.

The invention also provides a pharmaceutical pack or  
30   kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions

-48-

of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, the polypeptides of the present invention may be employed in conjunction with other therapeutic compounds.

10        V.    Polypeptide Production Methods

One embodiment of the present invention relates to the substantially purified polypeptide encoded by a mFLINT gene or a mFLINT gene.

15        Skilled artisans will recognize that the polypeptides of the present invention can be synthesized by a number of different methods, such as chemical methods well known in the art, including solid phase peptide synthesis or recombinant methods. Both methods are described in U.S. Patent 4,617,149.

20        The principles of solid phase chemical synthesis of polypeptides are well known in the art and may be found in general texts in the area. For example, see H. Dugas and C. Penney, BIOORGANIC CHEMISTRY (1981) Springer-Verlag, New York, 54-92. For example, peptides may be synthesized by solid-phase methodology utilizing an Applied Biosystems 430A peptide synthesizer (Applied Biosystems, Foster City, CA) and synthesis cycles supplied by Applied Biosystems.

25        The polypeptides of the present invention can also be produced by recombinant DNA methods using the cloned FLINT gene. Recombinant methods are preferred if a high yield is

30

-49-

desired. Expression of the cloned gene can be carried out in a variety of suitable host cells, well known to those skilled in the art. For this purpose, the FLINT gene is introduced into a host cell by any suitable means, well known to those skilled in the art. While chromosomal integration of the cloned gene is within the scope of the present invention, it is preferred that the gene be cloned into a suitable extra-chromosomally maintained expression vector so that the coding region of the FLINT gene is operably-linked to a constitutive or inducible promoter.

The basic steps in the recombinant production of the FLINT polypeptide are:

- a) constructing a natural, synthetic or semi-synthetic DNA encoding FLINT polypeptide;
- 15      b) integrating said DNA into an expression vector in a manner suitable for expressing the FLINT polypeptide, either alone or as a fusion polypeptide;
- c) transforming or otherwise introducing said vector into an appropriate eukaryotic or prokaryotic host cell forming a recombinant host cell;
- 20      d) culturing said recombinant host cell in a manner to express the FLINT polypeptide; and
- e) recovering and substantially purifying the FLINT polypeptide by any suitable means well known to those skilled in the art.
- 25

#### **1. Expressing Recombinant FLINT Polypeptide in Prokaryotic and Eukaryotic Host Cells**

Prokaryotes may be employed in the production of recombinant FLINT polypeptide. For example, the *Escherichia coli* K12 strain 294 (ATCC No. 31446) is particularly useful for the prokaryotic expression of foreign polypeptides.

-50-

Other strains of *E. coli*, bacilli such as *Bacillus subtilis*, enterobacteriaceae such as *Salmonella typhimurium* or *Serratia marcescans*, various *Pseudomonas* species and other bacteria, such as *Streptomyces*, may also be employed as host  
5 cells in the cloning and expression of the recombinant polypeptides of this invention.

Promoter sequences suitable for driving the expression of genes in prokaryotes include -lactamase [e.g. vector pGX2907, ATCC 39344, contains a replicon and -lactamase  
10 gene], lactose systems [Chang et al., *Nature* (London), 275:615 (1978); Goeddel et al., *Nature* (London), 281:544 (1979)], alkaline phosphatase, and the tryptophan (*trp*) promoter system [vector pATH1 (ATCC 37695)], which is designed to facilitate expression of an open reading frame  
15 as a *trpE* fusion polypeptide under the control of the *trp* promoter. Hybrid promoters such as the *tac* promoter (isolatable from plasmid pDR540, ATCC-37282) are also suitable. Still other bacterial promoters, whose nucleotide sequences are generally known, may be ligated to DNA  
20 encoding the polypeptide of the instant invention, using linkers or adapters to supply any required restriction sites. Promoters for use in bacterial systems also will contain a Shine-Dalgarno sequence operably-linked to the DNA encoding the desired polypeptides. These examples are  
25 illustrative rather than limiting.

The polypeptides of this invention may be synthesized either by direct expression or as a fusion polypeptide comprising the polypeptide of interest as a translational fusion with another polypeptide or peptide that may be  
30 removed by enzymatic or chemical cleavage. It often is observed in the production of certain peptides in

-51-

recombinant systems that expression as a fusion polypeptide prolongs the lifespan, increases the yield of the desired peptide, or provides a convenient means of purifying the polypeptide. This is particularly relevant when expressing mammalian polypeptides in prokaryotic hosts. A variety of peptidases (e.g. enterokinase and thrombin) which cleave a polypeptide at specific sites or digest the peptides from the amino or carboxy termini (e.g. diaminopeptidase) of the peptide chain are known. Furthermore, particular chemicals (e.g. cyanogen bromide) will cleave a polypeptide chain at specific sites. The skilled artisan will appreciate the modifications necessary to the amino acid sequence (and synthetic or semi-synthetic coding sequence if recombinant means are employed) to incorporate site-specific internal cleavage sites. For instance, see P. Carter, "Site Specific Proteolysis of Fusion Polypeptides", Chapter 13, in PROTEIN PURIFICATION: FROM MOLECULAR MECHANISMS TO LARGE SCALE PROCESSES, American Chemical Society, Washington, D.C. (1990).

In addition to prokaryotes, a variety of amphibian expression systems such as frog oocytes, and mammalian cell systems can be used. The choice of a particular host cell depends to some extent on the particular expression vector used. Exemplary mammalian host cells suitable for use in the present invention include HepG-2 (ATCC HB 8065), CV-1 (ATCC CCL 70), LC-MK<sub>2</sub> (ATCC CCL 7.1), 3T3 (ATCC CCL 92), CHO-K1 (ATCC CCL 61), HeLa (ATCC CCL 2), RPMI8226 (ATCC CCL 155), H4IIEC3 (ATCC CCL 1600), C127I (ATCC CCL 1616), HS-Sultan (ATCC CCL 1484), and BHK-21 (ATCC CCL 10), for example.

-52-

A wide variety of vectors are suitable for transforming mammalian host cells. For example, the pSV2-type vectors comprise segments of the simian virus 40 (SV40) genome required for transcription and polyadenylation. A large  
5 number of plasmid pSV2-type vectors have been constructed, such as pSV2-gpt, pSV2-neo, pSV2-dhfr, pSV2-hyg, and pSV2-globin, in which the SV40 promoter drives transcription of an inserted gene. These vectors are widely available from sources such as the American Type Culture Collection (ATCC),  
10 12301 Parklawn Drive, Rockville, Maryland, 20852, or the National Center for Agricultural Utilization Research, 1815 North University Street, Peoria, Illinois 61604-39999.

Promoters suitable for expression in mammalian cells include the SV40 late promoter, promoters from eukaryotic  
15 genes, such as, for example, the estrogen-inducible chicken ovalbumin gene, the interferon genes, the glucocorticoid-inducible tyrosine aminotransferase gene, the thymidine kinase gene promoter, and the promoters of the major early and late adenovirus genes.

20 Plasmid pRSVcat (ATCC 37152) comprises portions of a long terminal repeat of the Rous Sarcoma virus, a virus known to infect chickens and other host cells. This long terminal repeat contains a promoter which is suitable for use in the vectors of this invention. H. Gorman et al.,  
25 *Proc. Nat. Acad. Sci. (USA)*, 79, 6777 (1982). The plasmid pMSVi (NRRL B-15929) comprises the long terminal repeats of the Murine Sarcoma virus, a virus known to infect mouse and other host cells. The mouse metallothionein promoter has also been well characterized for use in eukaryotic host  
30 cells and is suitable for use in the present invention. This promoter is present in the plasmid pdBPV-MMTneo (ATCC

-53-

37224) which can serve as the starting material for the construction of other plasmids of the present invention.

Transfection of mammalian cells with vectors can be performed by a plurality of well known processes including, but not limited to, protoplast fusion, calcium phosphate co-precipitation, electroporation and the like. See, e.g., Maniatis et al., *supra*.

Some viruses also make appropriate vectors. Examples include the adenoviruses, the adeno-associated viruses, the vaccinia virus, the herpes viruses, the baculoviruses, and the rous sarcoma virus, as described in U.S. Patent 4,775,624, incorporated herein by reference. For example, the baculovirus pFastBac-1 (GIBCO/BRL) can be used to infect a suitable host cell, such as SF9, to produce recombinant protein.

Eukaryotic microorganisms such as yeast and other fungi are also suitable host cells. The yeast *Saccharomyces cerevisiae* is the preferred eukaryotic microorganism. Other yeasts such as *Kluyveromyces lactis* and *Pichia pastoris* are also suitable. For expression in *Saccharomyces*, the plasmid YRp7 (ATCC-40053), for example, may be used. See, e.g., L. Stinchcomb et al., *Nature*, 282, 39 (1979); J. Kingsman et al., *Gene*, 7, 141 (1979); S. Tschemper et al., *Gene*, 10, 157 (1980). Plasmid YRp7 contains the TRP1 gene which provides a selectable marker for use in a *trp1* auxotrophic mutant.

#### Purification of Recombinantly-Produced FLINT Polypeptide

An expression vector carrying the cloned FLINT gene is transformed or transfected into a suitable host cell using standard methods. Cells that contain the vector are propagated under conditions suitable for expression of the

-54-

recombinant FLINT polypeptide. For example, if the recombinant gene has been placed under the control of an inducible promoter, suitable growth conditions would incorporate the appropriate inducer. The recombinantly-produced polypeptide may be purified from cellular extracts of transformed cells by any suitable means.

In a preferred process for polypeptide purification, the FLINT gene is modified at the 5' end to incorporate several histidine residues at the amino terminus of the FLINT polypeptide. This "histidine tag" enables a single-step polypeptide purification method referred to as "immobilized metal ion affinity chromatography" (IMAC), essentially as described in U.S. Patent 4,569,794, which hereby is incorporated by reference. The IMAC method enables rapid isolation of substantially pure recombinant FLINT polypeptide starting from a crude extract of cells that express a modified recombinant polypeptide, as described above.

Other embodiments of the present invention comprise isolated nucleic acids that encode Figures 1, 2, 3, 4. Any of these nucleic acids may be produced by chemical synthetic methods. The synthesis of nucleic acids is well known in the art. See, e.g., E.L. Brown, R. Belagaje, M.J. Ryan, and H.G. Khorana, *Methods in Enzymology*, 68:109-151 (1979). Fragments of the DNA sequence corresponding to a FLINT or an mFLINT gene could be generated using a conventional DNA synthesizing apparatus, such as the Model 380A or 380B DNA synthesizers of Applied Biosystems, Inc. (850 Lincoln Center Drive, Foster City, CA 94404), using phosphoramidite chemistry, and thereafter ligating the fragments so as to reconstitute the entire gene. Alternatively,



-55-

phosphotriester chemistry may be employed to synthesize nucleic acids used in this invention. See, for example, Gait, M.J., ed. OLIGONUCLEOTIDE SYNTHESIS, A PRACTICAL APPROACH (1984).

5        In an alternative methodology, namely PCR, the DNA sequences disclosed and described herein, comprising, for example, Figure 1 can be produced from a plurality of starting materials. For example, starting with a cDNA preparation (e.g. cDNA library) derived from tissue that  
10        expresses the FLINT gene, suitable oligonucleotide primers complementary to Figure 1 or to any sub-region therein, are prepared as described in U.S. Patent No. 4,889,818. Other suitable protocols for the PCR are disclosed in PCR  
15        PROTOCOLS: A GUIDE TO METHOD AND APPLICATIONS, Ed. Michael A. Innis et al., Academic Press, Inc. (1990).

      The ribonucleic acids of the present invention may be prepared via polynucleotide synthetic methodology, discussed above, or they may be prepared enzymatically, for example, by using RNA polymerase to transcribe a FLINT DNA template.  
20        The most preferred systems for preparing the ribonucleic acids of the present invention employ the RNA polymerase from the bacteriophage T7 or the bacteriophage SP6. These RNA polymerases are highly specific, requiring the insertion of bacteriophage-specific sequences at the 5' end of the  
25        template to be transcribed. See Maniatis et al., *supra*. This invention also provides nucleic acids, RNA or DNA, that are complementary to Figures 1-4.

## 2. Vectors

      Another aspect of the present invention relates to  
30        recombinant DNA cloning vectors and expression vectors comprising the nucleic acids of the present invention.

-56-

The skilled artisan understands that choosing the most appropriate cloning vector or expression vector depends upon a number of factors including the availability of restriction enzyme sites, the type of host cell into which the vector is to be transfected or transformed, the purpose of the transfection or transformation (e.g., stable transformation as an extrachromosomal element, or integration into the host chromosome), the presence or absence of readily assayable or selectable markers (e.g., antibiotic resistance and metabolic markers of one type and another), and the number of copies of the gene desired in the host cell.

Vectors suitable to carry the nucleic acids of the present invention comprise RNA viruses, DNA viruses, lytic bacteriophages, lysogenic bacteriophages, stable bacteriophages, plasmids, viroids, and the like. The most preferred vectors are plasmids.

When preparing an expression vector the skilled artisan understands that there are many variables to be considered, for example, whether to use a constitutive or inducible promoter. The practitioner also understands that the amount of nucleic acid or polypeptide to be produced dictates, in part, the selection of the expression system. Regarding promoter sequences, inducible promoters are preferred because they enable high level, regulatable expression of an operably-linked gene. The skilled artisan will recognize a number of suitable promoters that respond to a variety of inducers, for example, carbon source, metal ions, and heat. Other relevant considerations regarding an expression vector include whether to include sequences for directing the localization of a recombinant polypeptide. For example, a

-57-

sequence encoding a signal peptide preceding the coding region of a gene is useful for directing the extra-cellular export of a resulting polypeptide.

The present invention also provides a method for constructing a recombinant host cell capable of expressing polypeptides comprising Figures 1-4. This method comprises transforming or otherwise introducing into a host cell a recombinant DNA vector that comprises an isolated DNA sequence as described in any of Figures 1 through 4.

The preferred host cell is any eukaryotic cell that can accommodate high level expression of an exogenously introduced gene or polypeptide, and that will incorporate said polypeptide into its membrane structure. Vectors for expression are those which comprise any of the sequences of Figures 1-4. Transformed host cells may be cultured under conditions well known to skilled artisans such that FLINT or mFLINT is expressed, thereby producing a recombinant FLINT or mFLINT polypeptide in the recombinant host cell.

The following examples more fully describe the present invention. Those skilled in the art will recognize that the particular reagents, equipment, and procedures described are merely illustrative and are not intended to limit the present invention in any manner.

#### EXAMPLE 1

##### RT-PCR Amplification of FLINT Gene from mRNA

A FLINT gene is isolated by reverse transcriptase PCR (RT-PCR) using conventional methods. Total RNA from a tissue that expresses the FLINT gene, for example, lung, is prepared using standard methods. First strand FLINT cDNA synthesis is achieved using a commercially available kit (SuperScript™ System; Life Technologies) by PCR in

-58-

conjunction with specific primers directed at any suitable region of Figures 1-4.

Amplification is carried out by adding to the first strand cDNA (dried under vacuum): 8  $\mu$ l of 10X synthesis buffer (200 mM Tris-HCl, pH 8.4; 500 mM KCl, 25 mM MgCl<sub>2</sub>, 1 ug/ $\mu$ l BSA); 68  $\mu$ l distilled water; 1  $\mu$ l each of a 10 uM solution of each primer; and 1  $\mu$ l Taq DNA polymerase (2 to 5 U/ $\mu$ l). The reaction is heated at 94° C for 5 minutes to denature the RNA/cDNA hybrid. Then, 15 to 30 cycles of PCR amplification are performed using any suitable thermal cycle apparatus. The amplified sample may be analyzed by agarose gel electrophoresis to check for an appropriately-sized fragment.

#### EXAMPLE 2

##### 15 Production of a Vector for Expressing FLINT in a Host Cell

An expression vector suitable for expressing FLINT or fragment thereof in a variety of prokaryotic host cells, such as *E. coli* is easily made. The vector contains an origin of replication (Ori), an ampicillin resistance gene (Amp) useful for selecting cells which have incorporated the vector following a transformation procedure, and further comprises the T7 promoter and T7 terminator sequences in operable linkage to a FLINT coding region. Plasmid pET11A (obtained from Novogen, Madison WI) is a suitable parent  
25 plasmid. pET11A is linearized by restriction with endonucleases NdeI and BamHI. Linearized pET11A is ligated to a DNA fragment bearing NdeI and BamHI sticky ends and comprising the coding region of the FLINT gene as disclosed by Figures 1-4.

-59-

The FLINT gene used in this construction may be slightly modified at the 5' end (amino terminus of encoded polypeptide) in order to simplify purification of the encoded polypeptide product. For this purpose, an oligonucleotide encoding 8 histidine residues is inserted after the ATG start codon. Placement of the histidine residues at the amino terminus of the encoded polypeptide serves to enable the IMAC one-step polypeptide purification procedure.

### EXAMPLE 3

#### Recombinant Expression and Purification of FLINT Polypeptide

An expression vector that carries an open reading frame (ORF) encoding FLINT or fragment thereof and which ORF is operably-linked to an expression promoter is transformed into *E. coli* BL21 (DE3) (*hsdS gal cIts857 ind1Sam7nin5lacUV5-T7gene 1*) using standard methods. Transformants, selected for resistance to ampicillin, are chosen at random and tested for the presence of the vector by agarose gel electrophoresis using quick plasmid preparations. Colonies which contain the vector are grown in L broth and the polypeptide product encoded by the vector-borne ORF is purified by immobilized metal ion affinity chromatography (IMAC), essentially as described in US Patent 4,569,794.

Briefly, the IMAC column is prepared as follows. A metal-free chelating resin (e.g., Sepharose 6B IDA, Pharmacia) is washed in distilled water to remove preservative substances and infused with a suitable metal ion [e.g., Ni(II), Co(II), or Cu(II)] by adding a 50mM metal chloride or metal sulfate aqueous solution until about 75% of the interstitial spaces of the resin are saturated with

-60-

colored metal ion. The column is then ready to receive a crude cellular extract containing the recombinant polypeptide product.

After removing unbound polypeptides and other materials by washing the column with any suitable buffer, pH 7.5, the bound polypeptide is eluted in any suitable buffer at pH 4.3, or preferably with an imidazole-containing buffer at pH 7.5.

#### EXAMPLE 4

##### Tissue Distribution of FLINT mRNA

The presence of FLINT mRNA in a variety of human tissues was analyzed by Northern analysis. Total RNA from different tissues or cultured cells was isolated by a standard guanidine chloride/phenol extraction method, and poly-A<sup>+</sup> RNA was isolated using oligo(dT)-cellulose type 7 (Pharmacia). Electrophoresis of RNA samples was carried out in formaldehyde followed by capillary transfer to Zeta-Probe™ nylon membranes (Bio-Rad, Hercules, Calif.). Figure 1 was the template for generating probes using a MultiPrime™ random priming kit (Amersham, Arlington Heights, Ill.). The efficiency of the labeling reaction was approximately  $4 \times 10^{10}$  cpm incorporated per  $\mu\text{g}$  of template. The hybridization buffer contained 0.5M sodium phosphate, 7% SDS (wt/vol), 1% BSA (wt/vol), and 1 mM EDTA. Prehybridization was carried out in hybridization buffer at 65° C for 2 h and <sup>32</sup>P-labeled probe was added and incubation continued overnight. The filters were washed in Buffer A (40 mM sodium phosphate pH 7.2, 5% SDS [wt/vol], 0.5% BSA [wt/vol], and 1 mM EDTA) at 65° C for 1 h, and then in Buffer B (40 mM sodium phosphate, pH 7.2, 1% SDS [wt/vol], and 1 mM EDTA) at 65° C for 20

-61-

minutes. The filters were air-dried and exposed to Kodak X-OMAT AR film at -80° C with an intensifying screen.

The results showed that FLINT mRNA was present in numerous tissues, including stomach, spinal cord, lymph  
5 node, trachea, spleen, colon and lung.

#### EXAMPLE 5

##### Production of an Antibody to a Polypeptide

Substantially pure polypeptide or fragment thereof is isolated from transfected or transformed cells using any of  
10 the well known methods in the art, or by a method specifically disclosed herein. Concentration of polypeptide in a final preparation is adjusted, for example, by filtration through an Amicon filter device such that the level is about 1 to 5 ug/ml. Monoclonal or polyclonal  
15 antibody can be prepared as follows.

Monoclonal antibody can be prepared from murine hybridomas according to the method of Kohler and Milstein (Nature, 256, 495, 1975), or a modified method thereof. Briefly, a mouse is repetitively inoculated with a few  
20 micrograms of the polypeptide or fragment thereof, or fusion peptide thereof, over a period of a few weeks. The mouse is then sacrificed and the antibody producing cells of the spleen isolated. The spleen cells are fused by means of polyethylene glycol with mouse myeloma cells. Fused cells  
25 that produce antibody are identified by any suitable immunoassay, for example, ELISA, as described in E. Engvall, Meth. Enzymol., 70, 419, 1980.

Polyclonal antiserum can be prepared by well known methods, as described, for example, by J. Vaitukaitis  
30 et.al., Clin. Endocrinol. Metab. 33, 988 (1971), that involve immunizing suitable animals with the polypeptides,

-62-

fragments thereof, or fusion polypeptides thereof, disclosed herein. Small doses (e.g., nanogram amounts) of antigen administered at multiple intradermal sites appears to be the most reliable method.

5

**EXAMPLE 6****Construction of Mammalian FLINT-Flag Expression Vector**

To facilitate confirmation of FLINT expression (without the use of antibodies), a bicistronic expression vector (pIG1-FLINTF) was constructed by insertion of an "internal  
10 ribosome entry site"/enhanced green fluorescent polypeptide (IRES/eGFP) PCR fragment into the mammalian expression vector pGTD (Gerlitz, B. et al., 1993, Biochemical Journal 295:131). This new vector, designated pIG1, contains the following sequence landmarks: the Ela-responsive GBMT  
15 promoter (D. T. Berg et al., 1993 BioTechniques 14:972; D.T. Berg et al., 1992 Nucleic Acids Research 20:5485); a unique BclI cDNA cloning site; the IRES sequence from encephalomyocarditis virus (EMCV); the eGFP (Clontech) coding sequence (Cormack, et al., 1996 Gene 173:33); the  
20 SV40 small "t" antigen splice site/poly-adenylation sequences; the SV40 early promoter and origin of replication; the murine dihydrofolate reductase (dhfr) coding sequence; and the pBR322 ampicillin resistance marker/origin of replication. The resultant protein had the  
25 Flag sequence attached at the C-terminal end, yielding:  
[FLINT]-DYKDDDDK.

Based upon the human FLINT sequence, the following primers were synthesized: 5'-

TAGGGCTGATCAAGGATGGGCTTCTGGACTTGGGCGGCCCC

30 TCCGCAGGCGGACCGGGG-3'; and 5'-AGGGGGGCGGCCGCTGATCATCACTT



-63-

GTCGTCGTCGTCCTTGTAGTCGTGCA CAGGGAGGAAGCGC - 3'. The latter, reverse primer contains the Flag epitope sequence (positions 24-47, double underline) (Micele, R.M. et al., 1994 J. Immunol. Methods 167:279). These primers were then used to

5 PCR amplify the FLINT cDNA. The resultant 1.3 Kb PCR product was then digested with *BclI* (restriction sites incorporated into primers, underlined above) and ligated into the unique *BclI* site of pIG1 to generate the plasmid pIG1-FLINTF. The human FLINT cDNA orientation and

10 nucleotide sequence were confirmed by restriction digest and double stranded sequencing of the insert.

#### EXAMPLE 7

##### Construction of Mammalian FLINT-non-Flag Expression Vector

In order to generate a non-Flagged expression vector

15 (pIG1-FLINT), the 24-base DNA sequence encoding the eight amino acid FLAG epitope was deleted from the pIG1-FLINTF construct using the Quick Change mutagenesis kit (Stratagene). A 35-base primer, and its complement, with identity to the 19-base sequences flanking the FLAG sequence

20 was synthesized and used for PCR amplification using pIG1-FLINTF the plasmid as template. The PCR reaction mixture was digested with *DpnI* restriction endonuclease to eliminate the parental DNA, and the PCR product was transformed into Epicurean XLI-blue E. coli cells. Sixteen ampicillin-

25 resistant transformants were picked and the plasmid DNA was analyzed by restriction digestion. Ten of the 16 gave results compatible with deletion of the 24-base sequence. Precise deletion of the 24-base sequence was confirmed by DNA sequencing of pIG1-FLINT. The nucleotide sequence of

30 FLINT in this plasmid is shown in Figure 1.

#### EXAMPLE 8

-64-

Isolation of a high-producing FLINT clone from AV12 RGT18  
transfectants

The recombinant plasmid carrying the FLINT gene (pIG1-FLINT) encodes resistance to methotrexate. In addition, the construct contains a gene encoding a fluorescent polypeptide, GFP, on the same transcript and immediately 3' to the FLINT gene. Since high level expression of GFP would require a high level of expression of the FLINT-GFP mRNA, highly fluorescent clones would have a greater probability of producing high levels of FLINT. pIG1-FLINT and pIG1-FLINTF were used to transfect AV12 RGT18 cells. Cells resistant to 250 nM methotrexate were selected and pooled. The pool of resistant clones was subjected to fluorescence assisted cell sorting (FACS), and cells having fluorescence values in the top 5% of the population were sorted into a pool and as single cells. The high fluorescence pools were subjected to three successive sorting cycles. Pools and individual clones from the second and third cycles were analyzed for FLINT production by SDS-PAGE. Pools or clones expressing FLINT at the highest level judged from Coomassie staining were used for scale-up and FLINT purification. The amino acid sequence of the mature FLINT protein that was produced using this plasmid is shown in Figure 3, called mFLINT.

25

EXAMPLE 9

Large Scale mFLINT Polypeptide Purification

Large scale production of mFLINT was carried out by first growing stable clones stable pIG1-FLINT-containing AV12 RGT 18 cells in several 10 liter spinners. After reaching confluency, cells were further incubated for 2-3

30

-65-

more days to secrete maximum amount of FLINT into media. Media containing mFLINT was adjusted to 0.1% CHAPS and concentrated in an Amicon ProFlux M12 tangential filtration system to 350 ml. The concentrated media was centrifuged at 19,000 rpm (43,000 x g) for 15 minutes and passed over a SP-5PW TSK-GEL column (21.5 mm x 15 cm; TosoHaas) at a flow rate of 8 ml/min. The column was washed with buffer A (20 mM MOPS, 0.1% CHAPS, pH 6.5) until the absorbency (280 nm) returned to baseline and the bound polypeptides were eluted with a linear gradient from 0.1 M-0.3 M NaCl (in buffer A) developed over 85 min. Fractions containing mFLINT were pooled and passed over a (7.5 mm x 7.5 cm) Heparin-5PW TSK-GEL column equilibrated in buffer B (50 mM Tris, 0.1% CHAPS, 0.3 M NaCl, pH 7.0). The bound polypeptide was eluted with a linear gradient from 0.3 M-1.0 M NaCl (in buffer B) developed over 60 min. Fractions containing mFLINT were pooled and passed over a 1 cm x 15 cm Vydac C4 column equilibrated with 0.1% TFA/H<sub>2</sub>O. The bound mFLINT was eluted with a linear gradient from 0-100% CH<sub>3</sub>CN/0.1% TFA. Fractions containing mFLINT were analyzed by SDS-PAGE and found to be greater than 95% pure and were dialyzed against 8 mM NaPO<sub>4</sub>, 0.5 M NaCl, 10% glycerol, pH 7.4. The N-terminal sequence of mFLINT was confirmed on the purified polypeptide. Mass spectral analysis and Endoglycosidase-F digestion indicates that mFLINT is glycosylated.

## EXAMPLE 10

A. Construction of FLINT-Flag and FLINT

Expression Vectors Flag-tagged and native versions of human FLINT were expressed in baculovirus-infected cells in order to generate enough recombinant protein for study. The human FLINT cDNA was engineered for expression as follows. A pIG3

-66-

(a derivative of pIG1; Gerlitz, B and Grinnell, B. W., unpublished data) expression vector containing the cDNA encoding a FLAG-tagged version of FLINT was cleaved with XbaI and filled in to create blunt ends. The vector was then  
5 cleaved with SalI. The resulting 918-base pair fragment containing the coding region was gel-purified and ligated into the baculovirus vector pFastBac-1 (GIBCO/ BRL) that had been digested with BamHI, the ends blunted and subsequently digested with SalI generating the plasmid pBacOPG3Flag. This  
10 construct was designed to express a full-length molecule (including the 29 NH2-terminal amino acids, which constitute the signal peptide) and the FLAG tag at the COOH-terminus of the protein. The expression was under the control of the baculovirus polyhedrin promoter.

15

For the construction of a vector to express the native version of the protein, a 920-base pair XbaI/ HindIII cDNA fragment encoding the full-length of the protein previously subcloned into pCDNA3.1 (+/-) [purchased from Invitrogen]  
20 was cleaved with XbaI and HindIII. The fragment was gel-purified and ligated into the baculovirus vector pFastBac-1 (GIBCO/ BRL) that had been digested with XbaI and HindIII generating the plasmid pBacOPG3.

25

#### B. Generation of Baculoviruses and Protein

Production The two vectors, pBacOPG3Flag and pBacOPG3 were separately used to generate two recombinant baculoviruses (vBacOPG3Flag and vBacOPG3) as described by the vendor (GIBCO/ BRL). Each of the viruses was separately used to  
30 infect SF-9 cells for protein production. The recombinant proteins were measured in supernatants collected from the

-67-

infected SF-9 cells by Western-blot and Coomassie stain analyses.

EXAMPLE 11

FAS Ligand Binding Experiments

To detect mFLINT interaction with FasL

5

Dot blot experiment was performed to scan known TNF ligands that are commercially available TRAIL and FasL for interaction with mFLINT.

10 TRAIL (RnD Systems) and FasL (Kamiya Biomedical Company) were spotted on a nitrocellulose paper and incubated with purified mFLINT-Flag. mFLINT was washed away and binding mFLINT was detected using anti Flag antibody. Both OPG2Fc and mFLINT-Flag were overexpressed and purified  
15 according the examples above. The filter paper was subsequently blocked for 30 min using 5% nonfat milk in PBS in room temperature. The nitrocellulose paper was subsequently mixed with the cell lysate containing FasL-Myc, and further incubated on a rotator for 1 hour at room  
20 temperature. Secondary and tertiary incubations were performed with anti-myc antibody and anti-mouse IgG-HRP for 1 hour and 30 minutes respectively. The polypeptide containing myc epitope was detected by chemiluminescence on X-ray film which showed that mFLINT bound to FasL  
25 specifically. No appreciable binding was detected with TNF , TNF , TRAIL, CD40L or TRANCE.

First a baseline experiment was done for the Fas-FasLigand interaction in vitro. Unless otherwise indicated, all washing steps use TBST (Tris Buffer Saline with Tween 20  
30 from SIGMA) and were done 3 to 6 times.

-68-

mrecFas (100 ng) was adsorbed on to ELISA plate. Then the plate was is blocked by TBST plus 0.1% Gelatine. Thereafter, hFasLigand (Flag-tagged) was added at different concentrations with a maximum concentration of 300 ng going  
5 down to 1 ng on TBST plus a 0.1% solution containing 1 micrograms/ml of M2 Abs (antiflag antibodies purchased through Scientific Imaging System division of Kodak). After washing the plate 6 times, anti-mouse-Abs-HRP (3000 dilution, Bio-Rad) was added to the wells. After washings  
10 three times, visualization enzymatic reaction using ABTS as a substrate was performed. Unless otherwise noted, an ELISA reader commercialized by Molecular Devices Corp. (Menlo Park, California) was used.

The following data were collected:

15

FasL, ng	OD, 405nM
1	.1
5	.2
10	.3
50	.7
100	1.2
500	1.6

FLINT-Fas L binding may be confirmed and specifics of binding determined (e.g., kinetics, specificity, affinity, cooperativity, relative binding pattern, concentration)  
20 using real-time biomolecular interaction analysis. This technology confers the ability to study biomolecular interactions in real time, without labeling any of the interactants. In particular, it takes advantage of the

-69-

optical phenomenon surface plasmon resonance, and detection depends on changes in the mass concentration of macromolecules at the biospecific interface. Interactions are followed in real time, so that kinetic information is readily derived. In many cases, investigations can be performed without prior purification of components.

Measurements are accomplished using a BiaCore 2000 instrument. The instrument, accompanying chips, immobilization and maintenance kits and buffers are obtained from Biacore AB, Rapskatan 7, S-754 50 Uppsala, Sweden. FasL is obtained from Kamiya Biomedical Company, 910 Industry Drive, Seattle, WA 98188, Guanidine Isothiocyanate Solution from GibcoBRL, and mFLINT is prepared as in Examples 8 and 9.

Experiments using immobilized FasL loaded with a solution containing mFLINT. In this experiment, the  $K_D$  of the FasL-FLINT interaction was  $1.13 \times 10^{-7}$ , which is lower than the FasL binding Fas, which was found to be  $1.62 \times 10^{-7}$ . This is explained by the fact that FasL is a trimer, yet here FasL monomer was bound. Further experiments utilized bound FLINT with FasL in solution, allowing for trimer formation. The resultant  $K_D$  for FLINT-FasL in these experiments was 2 orders of magnitude better (i.e., the  $K_D$  was lower by two orders of magnitude). This observations indicate that FLINT should effectively compete with Fas for FasL binding, which likely explains the therapeutic benefit observed in experiments detailed below.

#### FLINT prevents Fas-FasLigand interaction

As above, mrecFas (100 ng) was adsorbed on to ELISA plate. Again the plate was blocked by TBST and

-70-

0.1% Gelatine. Thereafter, hFasLigand (Flag-tagged, 30 ng per each point) in the presence of different mFLINT concentrations (Maximum concentration 300 ng down to 1 ng) on TBST plus a 0.1% solution containing 1 microgram/ml of M2 Abs is added to each well. As before, after washing of the plate, anti-mouse-Abs-HRP (3000 dilution, Bio-Rad) was added to the wells. After washings, visualization enzymatic reaction using ABTS as a substrate. was performed. The data is shown in the following table.

10

FLINT, ng	OD, 405nm
1	0.36
5	0.36
10	0.36
50	0.28
100	0.18
500	0.06

FasLigand binds Fas and mFLINT with different affinities.

FLINT and Fas (100 ng of each) were adsorbed on to an ELISA plate. hFasLigand (Flag-tagged) was added at different concentrations to a maximum concentration of 300 ng down to 0.1 ng on TBST plus a 0.1% solution containing 1 microgram/ml of M2 Abs. After washing of the plate, anti-mouse-Abs-HRP (1:3000 dilution, Bio-Rad) was added to the wells. After washings, visualization enzymatic reaction using ABTS as a substrate was performed. The table below shows the data.



-71-

FasL, ng	FLINT OD	Fas OD, 405nm
0.1	0	0
0.5	0	0
1.0	.02	0
5.0	.04	.01
10	.12	.03
50	.28	.045
100	.78	.18

EXAMPLE 12Measuring the effect of mFLINT on anti-CD3 induced Jurkat apoptosis

5 Non-tissue treated 24 well plates (Decton Dickinson, Mansfield, MA) were coated with 0.5 ml of 1 ug/ml anti-CD3 (Farmingen) in PBS for 90 min at 37 °C. The plate was washed once with PBS. 1 ml of  $1 \times 10^6$  cell/ml was seated in each well with or without following treatment: 10  $\mu$ M DEVD-  
 10 cmk, 1 ug OPG2-Fc, 1 or 2 ug of mFLINT and 1 ug anti FasL Ab. mFLINT was made according to Examples 8 and 9.

Cells were incubated overnight at 37 °C incubator and cells were then stained by Annexin V and PI staining. Apoptosis was analyzed by flow cytometer (FACS). Cell  
 15 apoptosis was indicated by positive staining with Annexin V.

Control Jurkat	6.97
Jurkat + anti Fas	59.28
Jurkat + antiCD3	46.32
Jurkat + antiCD3 + DEVDcmk	30.80
Jurkat + antiCD3 + mFLINT (1ug)	27.77
Jurkat + antiCD3 + OPG2-Fc (1ug)	45.78

-72-

Jurkat + antiCD3 + mFLINT (2ug)	18.67
Jurkat + antiCD3 + antiFasL Ab	24.05

EXAMPLE 13

Measuring the effect of mFLINT on recombinant FasL induced  
Jurkat cells apoptosis

- 5 One milliliter of  $1 \times 10^6$  cell/ml was added into each well of 24 well tissue culture plate and treated with following reagents: soluble Fas L (200 ng), Fas L plus 1 ug mFLINT, Fas L plus 1 ug OPG2-Fc, Trail (200 ng), Trail plus 1 ug mFLINT. Cells were incubated overnight at 37 °C and
- 10 then stained with Annexin V and PI. Cell apoptosis was analyzed by flow cytometer (FACS). mFLINT was made according to Examples 8 and 9.

Control Jurkat	3.23
Jurkat + FasL (200ng/ml)	67.39
Jurkat + FasL (200ng/ml) + anti FasL Ab (1 ug)	3.3
Jurkat + FasL (200ng/ml) + mFLINT (1 ug)	3.32
Jurkat + FasL (200ng/ml) + mFLINT (1 ug)	4.6
Jurkat + FasL (200ng/ml) + OPG2 (1ug)	70.58
Jurkat + FasL (200ng/ml) + OPG2 (1ug)	69.58
Jurkat + TRAIL (200ng/ml)	17.47
Jurkat + TRAIL (200ng/ml)	17.43

EXAMPLE 14Measuring the effect of mFLINT in a dose-dependent manner on  
anti-CD3 induced Jurkat apoptosis

The same steps for plate coating and cell treatment set  
5 out in Example 13 were followed except a different amount of  
mFLINT was added into each well. mFLINT was made according  
to Examples 8 and 9. The following table indicates the  
amounts added:

Jurkat cells (Control)	5.33
Jurkat cells + anti CD3	27.49
Jurkat cells + anti CD3 + anti FasL neutralization Ab	12.74
Jurkat cells + anti CD3 + OPG2-Fc 4ug	26.24
Jurkat cells + anti CD3 + mFLINT/PG3 3000ng	14.68
Jurkat cells + anti CD3 + mFLINT 2000ng	17.02
Jurkat cells + anti CD3 + mFLINT 1000ng	24.29
Jurkat cells + anti CD3 + mFLINT 500ng	27.48
Jurkat cells + anti CD3 + mFLINT 250ng	28.93
Jurkat cells + anti CD3 + mFLINT 125ng	29.4
Jurkat cells + anti CD3 + mFLINT 62.5ng	28.99
Jurkat cells + anti CD3 + mFLINT	28.21

31.25ng	
Jurkat cells + anti CD3 + mFLINT	28.80
15.625ng	

### EXAMPLE 15

Measuring the effect of human mFLINT on murine FasL-mediated apoptosis using mouse T cell hybridoma cells (LTT cells)

5 (Annexin V assay)

FLINT was made according to Examples 8 and 9.

LTT, 2, 14, 11 cells (LTT cells), see Glasebrook, *Eur. J. Immunol.* 17: 1561-65 (1987), were used in this Annexin V assay. On the first day, a 96 well plate was coated with anti-CD3 (2C11) at a serial dilutions. On the second day, 100,000 LTT cells were added, in 50  $\mu$ l of medium per well, along with 50  $\mu$ l of medium to the control wells and 50  $\mu$ l of medium containing:

Group 1. solubleFas (sFas) (FasFc, mouse), with a final  
15 concentration of 1 µg/ml

Group 2. mFLINT (human), with a final concentration of 1 ug/ml

Group 3. anti-FasL (mouse), with a final concentration of 1 ug/ml.

20            These then were incubated overnight, at 37° C, in 5%  
              CO<sub>2</sub>.

On the following day (Day 3), the cells were collected from each well, washed and labeled with Annexin V and PI, following Flow cytometry analysis.

25

-75-

Anti- CD3	Contr ol  % Annexin V Positive	SFas % Annexin V Positive	FLINT % Annexin V Positive	Anti- FasL % Annexin V Positive
1	96.42	56.66	34.41	53.46
0.33	94.48	47.28	35.89	39.21
0.11	91.27	41.08	33.24	32.54
0	19.74	23.14	26.47	17.54

**EXAMPLE 16**

Measuring the effect of human mFLINT on murine FasL-mediated apoptosis using mouse T cell hybridoma cells (LTT cells

5 (Cytotoxicity Assay)

The same steps as in Example 16 were followed on Day 1 and Day 2. On Day 3 20ul of MTS solution (Promega) was added to the cells which were then incubated at 37°C for 2 hours. Using a plate reader, the absorbances at 490nm  
10 wavelength were collected.

Anti- CD3 conc. (µg/ml)	s-Fas (1 µg/ml))	FLINT (1 µg/ml)	Anti- FasL	Contr ol
0	1.774	1.691	2.01	1.534
0.0014	1.968	1.923	2.134	1.614
0.004	1.929	1.982	2.147	1.653
0.012	1.779	2.006	2.108	1.284
0.037	1.777	2.006	1.988	0.834
0.11	1.638	1.874	1.956	0.733
0.33	1.624	1.671	1.978	0.648

-76-

1	1.459	1.581	1.887	0.664
---	-------	-------	-------	-------

EXAMPLE 17LIGHT Binding experiments

To confirm the dot blot binding between LIGHT and  
5 mFLINT, 293 cells were transiently transfected with LIGHT  
expression construct overnight. On the next day, cells were  
detached and incubated with mFLINT-Flag on ice. The Flag  
epitope was subsequently detected by anti-Flag conjugated  
with fluorochrome and the cells population that shows  
10 specific binding with mFLINT was detected by flow cytometer.  
As a control we used vector transfected cells. To make sure  
that the binding was specific, a competition assay with 10  
fold excess of untagged mFLINT was performed.

More particularly, 6 well dishes of cells were  
15 transfected as above. Both vector and m-LIGHT expressing  
cells were provided. Cells were detached from the plates by  
vigorous pipetting with a P100 Pipettor. Then, in PBS/BSA,  
0.1%, cells were exposed to one of the following  
combinations, set forth in a-d.

- 20 a. GST/flag, mFLINT/flag, or HVEM at 20 nM;
- b. FLINT/flag at 20 nM + GST/flag or mFLINT or HVEM at  
200 nM;
- c. FLINT/flag at 20 nM + HVEM + anti-human FAS ligand  
at 200 nM;
- 25 d. anti-human FAS ligand-biotin at 1 g/ml

Cells were incubated on ice for 30 minutes and washed  
with PBS/BSA, 0.1%. Cells then were exposed either to anti-  
human IgG-biotin, 1 g/ml (for detection of HVEM), or to M2-

-77-

biotin, 2 µg/ml (for detection of flag conjugates). Cells were incubated on ice for 30 minutes and washed with PBS/BSA, 0.1%. Thereafter, cells were exposed to streptavidin Alexa 488 (SIGMA) at a 1:1000 dilution. Again, 5 cells were incubated on ice for 30 minutes and washed with PBS/BSA, 0.1%. Cells were analyzed using a FACSORT flow cytometer (Decton Dickinson) to determine binding.

The cell surface binding assay using flow cytometer confirmed that peaks shifted only when LIGHT expressing 10 cells were stained with mFLINT-Flag was used (data not shown). There was no shift in the control cells when stained with mFLINT-Flag. The shifted peak was completely reversed to baseline peak when the cells were preincubated with 10 fold excess of non tagged mFLINT thereby 15 preoccupying all the binding sites for mFLINT-Flag.

#### EXAMPLE 18

##### In vivo testing of mFLINT for treatment of Liver Damage

Using mice, a model of liver damage was induced using a 20 modification of the methods set out in Tsuji H., et al, 1997, Infection and Immunity, 65(5):1892-1898. mFLINT was made according to Examples 8 and 9.

Specifically, the activity of the polypeptides of the present invention against acute inflammation and apoptosis 25 was determined using the following procedure. Briefly, BALB/c mice (Harlan) per each experimental group were given intravenous injections (the lateral tail vein) of 6 mg of D(+)-Galactosamine (Sigma, 39F-0539) in 100 µl of PBS (GIBCO-BRL) and 3 µg of Lipopolysaccharide B *E.coli* 026:B6 30 (LPS) (Difco, 3920-25-2) in 100 µl of PBS. The LPS was

-78-

administered, via i.v. injection, 5 minutes after the galactosamine, which was administered i.v. After LPS challenge, the animals were injected intraperitoneally with mFLINT (200 µg), Hamster IgG (500 µg, Cappel, 30926), mAb  
5 against murine TNF, TN3-19.12 (500 µg, Sheehan K.C.F. et al J. Immunol. 1989. 142: 3884), and Anti-mouse Fas Ligand (500 µg, PharMingen, MO24301) at 0, 2, 4, 6 hour-point respectively. The survival rates of the mice were determined 24 and 48 hours after LPS injection.

10 In mice challenged with 3 µg of LPS after IP injection of 200 µg mFLINT polypeptide (FLINT) had a positive effect on animal survival. When mFLINT was administered 2 hours post-challenge, 100% of the animals survived; at 4 hours post- challenge, 73% of the animals survived; at 6 hours  
15 post-challenge, 60% of the animals survived. In contrast, administration of anti-TNF $\alpha$  at 4 hours post-challenge protected only 10% of the animals.

Figure 5 compares the effects of administering 200 µg mFLINT, i.v., and other molecules before and after challenge  
20 with LPS/GalN. When administered 2 hours before LPS/GalN, both anti-TNF and mFLINT treatment resulted in 100% survival of mice. 80% of mice treated with anti-FasL survived, and about 30% of mice treated with IgG survived.

In contrast, when anti-TNF was administered 4 hours  
25 after LPS/GalN, less than 10% of the animals survived at 48 hours. Surprisingly, 80% of the animals survived when treated with mFLINT, at 4 hours after LPS/GalN treatment. Administration of anti-FasL at 4 hours after LPS/GalN treatment also resulted in 80% survival. IgG had  
30 essentially no effect on survival, with only about 2% of the



-79-

animals surviving. Therefore, mFLINT is effective at treating late-stage disease.

Figure 6 shows that when 400 ug of mFLINT was administered i.p. to mice, 2 hours before challenge with LPS and GalN, 100% of the animals were alive 48 hours after LPS/GalN dosing. If 400 ug of anti-TNF was given i.p. two hours before LPS/GalN challenge, about 95% of the animals survived to 48 hours. In contrast, only 20% of those animals not treated with mFLINT survived 48 hours. Administration of IgG two hours before LPS/GalN only increased the survival rate to 30%.

Figure 6 shows that lowering the dose of mFLINT to 50 ug still had a protective effect. 48 hours after LPS/GalN administration, about 70% of the animals survived to 48 hours, compared with 0% survival of non-treated animals.

Figure 7 shows that the effect of 100 g mFLINT on survival is the same, whether given ip or iv, 12 or 2 hours before LPS/GalN administration. When mFLINT is given 4 hours after LPS/GalN administration, iv administration results in 100% survival, and IP administration results in 80% survival. This Figure also shows the dose/response relationship of the 4-hour post-LPS/GalN dose, given iv. 50 ug gave 80% survival, 10 ug gave 40% survival, 5 ug gave 20% survival and 1 ug gave only about 2% survival.

25

#### EXAMPLE 19

This example demonstrates the usefulness of mFLINT in the treatment of cerebral ischemia, which is clinically relevant in stroke, head trauma and other similar disorders.

Adult male gerbils (70 to 80 g body weight, Charles River Laboratories, Wilmington, MA) were anesthetized by

30

-80-

i.p. injections of sodium pentobarbital (Nembutal) 40 mg/kg, and additional i.p. injections of 10 mg/kg when necessary to maintain a surgical plane of anesthesia. Animals were placed on a thermostatically controlled heating blanket to  
5 maintain body temperature at 37 °C. The ventral surface of the neck was exposed, the fur shaved, and the skin cleaned with 2% iodine solution.

After the pre-surgical preparation, a midline incision was made, and the skin opened. The sternohyoid muscles  
10 were divided to exposed and isolate the common carotid arteries (CCA) for clamping. Sterilized aneurysm clips (blade with 0.15 mm, closing force ~10 gm) was secured by means of a sterilized clip applier on both left and right CCA for 5 minutes. The clamps were then removed and the  
15 patency of the arteries checked visually. The wound in the neck was closed by surgical suture.

Immediately following the cerebral ischemia procedure and while the gerbil was still unconscious, the fur on the dorsal surface of the head was shaved and the skin cleaned  
20 with 2% iodine solution. Under surgical anesthesia, the gerbil's head was secured in a stable position by means of a stereotaxic apparatus (SA) and a midline incision was made to expose the skull. At a position 1 mm lateral and 1 mm posterior to the bregma, as guided by the vernier scale of  
25 the SA, the skull was thinned by a dental drill equipped with a drill bit of 0.5 mm in diameter. The thinned area was punctured with a microsyringe equipped with a 27-gauge blunt needle inserted 3 mm deep for a bolus injection of 5 ul (0.63 mg/ml ) of mFLINT in phosphate buffer saline (PBS).  
30 mFLINT was made according to Examples 8 and 9.

-81-

After the bolus injection, the syringe needle was exchanged for an infusion cannula [3 mm in length] of a brain infusion assembly connected to an Alzet osmotic pump (Alza Corp., Palo Alto, CA) which reservoir was placed under the skin on the shoulder of the gerbil. The infusion cannula was anchored on the surface of the skull using dental cement. The wound was closed by surgical suture. The Alzet osmotic pump containing mFLINT solution (0.63 mg/ml) or PBS delivered continuously at a rate of 1  $\mu$ l/h for 3 days. Gerbils were allowed to survive for 5 days (the surgery day was taken as day zero).

On the fifth day of survival, the gerbils were sacrificed in a CO<sub>2</sub> chamber. Thoracotomy was performed for transcardiac perfusion of saline for 3 minutes and formaldehyde for 2 minutes. The brains were removed for histological processing following a standard procedure commonly adapted in the field. Coronal sections were obtained at approx. 1.7 mm posterior to the bregma. After staining with Cresyl violet, the sections were viewed under a microscope at 40x magnification for cell counter quantification of the intact hippocampal neurons along the dorsal CA1 regions (0.5 mm in length) of both hemispheres. Data were analyzed by Student t-Test and the Wilcoxon ranking test.

Results showed that mFLINT had a significant effect on neuronal survival compared to vehicle ( $p=0.0039$  in t-Test;  $p=0.0037$  in Wilcoxon Rank Sums) and was indistinguishable from normal controls.

-82-

EXAMPLE 20

TREATMENT GROUPS:	Number of Animals
CONTROLS	10
FLINT (50 µg/injection) iv, days 4-13	10
OPG2 (50 µg/injection) iv, days 4-13	<u>10</u>
	TOTAL: 30 mice

For the experiment, a single cell suspension of B16 melanoma cells was prepared from a brie of donor tumors.

5 Tumor cells ( $2 \times 10^6$ ) were implanted subcutaneously in a hind-leg of Male C57B1 mice from Taconic Farms on day 0. Treatment was initiated on day 4. mFLINT or OPG2 was administered by intravenous injection into a tail vein once daily on days 4 through 13 for a total of 10 injections.

10 mFLINT was made according to Examples 8 and 9. An overview of the protocol is shown above.

Tumor response was monitored by tumor volume measurements as determined by caliper measurements of two dimensions of the tumors on days 4, 8, 11, 17, 21, 25, 30

15 and 34. Tumor volume was calculated as a hemi-ellipsoid. The animals were weighed on the same schedule described above.

The control tumors reached a volume of  $500 \text{ mm}^3$  on day  $17.4 \pm 0.3$  and  $1000 \text{ mm}^3$  on day  $21.1 \pm 0.3$ . The tumors of

20 the animals treated with mFLINT reached a volume of  $500 \text{ mm}^3$

-83-

on day  $19.3 \pm 0.3$  and reached  $1000 \text{ mm}^3$  on day  $23.0 \pm 0.4$ . The tumors of the animals treated with OPG2 reached a volume of  $500 \text{ mm}^3$  on day  $185. \pm 0.4$  and reached  $1000 \text{ mm}^3$  on day  $22.2 \pm 0.4$ . Therefore, the tumor growth delay produced by mFLINT was 1.8 days and the tumor growth delay produced by OPG2 was 1.1 days. These data are shown graphically in Figure 8. As shown in Figure 8, tumor volume after 20 days was approximately  $730 \text{ mm}^3$  in mFLINT-treated mice, but the tumor volume of control mice was approximately  $1000 \text{ mm}^3$ . There was no indication of toxicity from administration of mFLINT or OPG2 as evidenced by weight loss in the animals.

#### EXAMPLE 21

Thirty NOD mice 6 weeks of age are purchased from Jackson Laboratories (Bar Harbor, Maine). The mice are housed three to a cage and given free access to food (Purina 5001) and water. After one week of acclimatization, the mice are bled by tail snip, and blood glucose and plasma insulin are measured. Blood glucose is measured by tail snip without anesthetic using a Precision G blood glucose analyzer (Medisense Inc., Bedford, MA). Plasma insulin is measured by a RIA kit (Linco Inc., St. Louis, MO.). The mice are then arbitrarily placed into three groups, control, mFLINT-injected, and OPG2-injected. Following assigning of the mice to groups, body weight and blood glucose are determined once weekly. Beginning at the onset of overt diabetes (blood glucose  $>200\text{mg\%}$ ; ~14 weeks of age) mice are injected once daily intraperitoneally with either mFLINT ( $50 \mu\text{g/day}$ ) or OPG2 ( $50 \mu\text{g/day}$ ) diluted in PBS. Control mice are injected with an equal volume of PBS. After 2 weeks of injection,

-84-

mice are anesthetized by inhalation anesthesia (carbon dioxide) and blood is removed by cardiac puncture for measurement of glucose and insulin. Additionally, the pancreas is harvested by the Animal Studies Support Team and  
5 fixed in zinc-formalin for 24 hours for subsequent histochemical analysis of  $\beta$  cell integrity (hematoxylin and eosin) and immunohistochemical staining for insulin (George Sanduskys laboratory). mFLINT is made according to Examples 8 and 9.

10

**EXAMPLE 22**

This example demonstrates that mFLINT inhibits apoptosis is an *in vitro* model that mimics ischemia reperfusion injury. These data indicate that mFLINT is useful in preventing and treating such injury.

15

Ventricular cardiomyocytes were isolated from the hearts of 1-3 day-old neonatal rats by trypsin digestion and non-myocytes were eliminated by pre-plating. Primary cultures were plated in special-coated 96-well plates in serum-free medium.

20

Hypoxia was induced by incubating cultures in glucose-free and oxygen-free (5% CO<sub>2</sub> and 95% N<sub>2</sub>) for 8 hours. Reperfusion was mimicked by a 16 hour incubation in glucose-containing medium under normal oxygen conditions. At the end of this incubation, cardiomyocyte apoptosis was measured  
25 by the cytoplasmic nucleosome-associated DNA ELISA method.

25

Test samples were incubated with mFLINT (2 g/ml or 5 g/ml) and control samples with Z-YVAD-fmk (50 M), a commercial caspase inhibitor. Duplicate experiments showed essentially complete inhibition of hypoxia/reperfusion-

30

induced apoptosis. mFLINT was made according to Examples 8 and 9.

-85-

To confirm these data and to query whether the observed inhibition was Fas-mediated, cardiomyocyte apoptosis was induced by incubation with soluble FasL, and apoptosis was measured as above. In these experiments, apoptosis was inhibited either by anti-Fas neutralizing antibody (1 g/ml) or mFLINT (10 g/ml). These experiments confirmed that mFLINT inhibits apoptosis, and indicated that it does so, at least in part, by inhibiting the FasL-Fas apoptotic pathway.

10

EXAMPLE 23Murine Osteoclast Differentiation Assay

The co-culture method of Takahashi et al. (Endocrinology 123:2600 1988) was modified as described in Galvin et al. (Endocrinology 137:2457 1996) and used to study the effects of various agents on osteoclast differentiation. mFLINT was made according to Examples 8 and 9.

Male Balb/C mice (4-8 weeks old) were euthanized with CO<sub>2</sub>, the femurs removed, and the marrow flushed out of the femurs with growth medium. Bone marrow cells were pelleted by centrifugation at 500 x g for 6 min. and resuspended in the growth medium (RPMI 1640 plus 5% heat inactivated fetal bovine-serum and 1% antibiotic-antimycotic solution). The marrow population ( $5 \times 10^4$  cells/cm<sup>2</sup>) was seeded in tissue culture dishes in which BALC cells (a stable cell line derived from neonatal mouse calvariae,  $1.5 \times 10^4$  cells/cm<sup>2</sup>) had been plated 2 h prior to addition of bone marrow. The cells were cultured for 7 days in a humidified incubator at 37°C with 5% CO<sub>2</sub> with medium changes on days 3 and 5. Cultures were treated with or without  $10^{-8}$ M 1,25-(OH)<sub>2</sub>D<sub>3</sub> on days 0, 3, and 5. In addition, the cells were treated with

-86-

or without secreted mFLINT protein purified from the conditioned medium of cells transfected with mFLINT gene (Figure 1). Following 7 days of culture, the cells in 24-well cluster dishes were fixed with formalin (3.7% for 10 min) and then stained for tartrate-resistant acid phosphatase (TRAP) using a modification of the method described by Graves, L and Jilka RL, J Cell Physiology 145:102 1990. The number of osteoclasts (TRAP-positive cells containing 3 or more nuclei) was quantitated. Results are reported in the following Table.

Table

FLINT (ng/ml)	Osteoclasts/well <sup>a</sup>
0.00	145.50 ± 7.33
0.01	40.50 ± 2.39*
0.10	65.50 ± 3.33*
1.00	97.50 ± 3.10*
10.00	170.17 ± 8.26
100.00	335.00 ± 8.90*

a - Each value represents the mean and standard error of 6 wells.

\*p<0.05 compared to control group

#### EXAMPLE 24

##### Porcine Osteoclast Differentiation Assay

Neonatal pigs (aged 1-5 days) were euthanized with CO<sub>2</sub>, the appendages were rinsed with 70% ethanol, the soft tissues were removed, and the humeri, radii, ulnae, femora, tibiae and fibulae were excised. The long bones were placed



-87-

in ice-cold calcium and magnesium-free Hank's balanced salt solution (CMF-HBSS, Gibco BRL) and cleaned of all soft tissues. The bones were split longitudinally and the endosteal surfaces were scraped to remove both the marrow and trabecular bone. The suspension of trabecular bone particles and marrow cells was agitated by vigorous shaking and passed through a 200 mm and then 100 mm sieve. Cells were centrifuged at 500 x g for 10 minutes at 4°C, the pellet was resuspended in CMF-HBSS, and then separated on a Ficol-Paque gradient (Pharmacia, Piscataway, NJ). The mononuclear cell fraction from the gradient was washed twice in CMF-HBSS and passed through a 35 mm sieve. The cells were suspended in growth medium consisting of  $\alpha$ -MEM (pH 7.2, which was modified to contain 8.3 mM NaHCO<sub>3</sub> (Gibco BRL, Grand Island, NY)), 10% heat-inactivated fetal bovine serum (FBS, Hyclone, Logan, UT) and 2% antibiotic/antimycotic solution (Gibco BRL, Grand Island, NY) and seeded onto tissue culture dishes at a density of  $1 \times 10^6$  cells/cm<sup>2</sup>. A typical marrow cell yield was between  $1-2 \times 10^9$  cells/animal, which varied with the size of the animal. The cells were incubated at 37°C in a humid incubator with 5% CO<sub>2</sub>. After 24-48 h, nonadherent cells were removed and seeded in either 24-well cluster dishes at a density of  $7.5 \times 10^5$  cells/cm<sup>2</sup> in growth medium which did or did not contain  $10^{-8}$  M 1,25-(OH)<sub>2</sub>D<sub>3</sub> (Biomol, Plymouth Meeting, PA) and mFLINT protein (obtained as in Examples 8 and 9). Cells were cultured for up to 10 days with medium changes every 48-72 h with growth medium that did or did not contain 1,25-(OH)<sub>2</sub>D<sub>3</sub> and mFLINT. Following 5 days of culture, the cells were fixed with formalin (3.7% for 10 min) and then stained for tartrate-resistant acid phosphatase (TRAP) as in

-88-

Example 6. The number of osteoclasts (TRAP-positive cells containing 3 or more nuclei) was quantitated. Results are reported in the following table.

5

Table

FLINT (ng/ml)	Osteoclasts/well <sup>a</sup>
0.00	214.83 + 14.22
0.01	68.83 + 6.28*
0.10	176.17 + 23.01
1.00	228.50 + 17.26
10.00	228.50 + 29.29
100.00	382.33 + 26.59*

a Each value represents the mean and standard error of 6 wells.

\*p<0.05 compared to control group

10

#### EXAMPLE 25

##### In vitro Treatment with mFLINT

FLINT was made according to Examples 8 and 9 and used in the following experiments.

15

A -Bone marrow cells from irradiated mice- This experiment was designed to determine whether mFLINT could improve the recovery of murine hematopoietic progenitor cells from irradiation, by forcing *in vitro* expansion with hematopoietic cytokines. Mice were injected with 3 mg of 5-flouro-uracil, i.p., and four days later, femurs were removed and flushed to isolate bone marrow (BM) cells. 1 x 10<sup>6</sup> BM cells were seeded in each well of a 24 well tray in triplicate in Iscoves modified dulbeccos media (IMDM)+ 10% FBS and were stimulated for 3 days with the cytokines stem

20

-89-

cell factor (SCF) and IL-6, in the presence or absence of mFLINT (1 ug/ml). After a 3-day *in vitro* expansion period,  $5 \times 10^3$  cells were plated out in triplicate for the CFU assay.

5 Bone marrow cells incubated with mFLINT had significantly increased numbers of CFUs compared to cells that were not treated with mFLINT. Figure 9 shows the CFU colonies/ $3 \times 10^6$  stimulated BM cells. These results suggest that mFLINT protected progenitor cells from apoptosis, and  
10 increased the number of viable cells that were able to form colonies. The Figure shows results for the following colony forming units- erythroid cells (CFU-E) (control - 15,000 colonies; mFLINT-treated - 33,000 colonies) ( $p < 0.003$ ) - granulocyte macrophage cells (CFU-GM) (control -- 8,000;  
15 mFLINT-treated - 15,000) ( $p < 0.039$ ); and the more primitive, granulocyte erythroid monocyte megakaryocyte cells (CFU-GEMM) (control - 750; mFLINT-treated-3,000) ( $p < 0.007$ ).  $p$  numbers show significant differences, as calculated by Student's T-test.

20 B- Bone marrow cells from mice treated with a chemotherapeutic agent -- This experiment was designed to demonstrate that mFLINT or anti-FasL antibody can improve the recovery of murine hematopoietic progenitor cells that have been exposed to a chemotherapeutic agent, following in  
25 *vitro* expansion with hematopoietic cytokines.  $2 \times 10^6$  bone marrow cells, from 5-FU treated mice, were seeded in each well of a 24 well tray in triplicate in IMDM media + 10% FBS and were stimulated for 3 days with SCF and IL-6, and with one of the following compounds:

- 30 1) mFLINT (1ug/ml)  
2) anti-FasL (1ug/ml)

-90-

- 3) FasL (0.15 ug/ml)
- 4) Control - no additions

Following a 3 day in vitro expansion period with  
5 cytokines, viable cells are counted and used to set up a CFU  
assay. If mFLINT or anti-FasL antibody protects progenitor  
cells from apoptosis, one would expect a greater number of  
colonies in these groups for the CFU assay. Results are  
shown in Fig. 10, which shows the bone marrow cell count  
10 (x1000). The values were: control cells: 3,300; FasL-  
treated cells: 3,000; anti-FasL-treated cells: 4,200;  
mFLINT-treated cells: 4,900.

C- CD34+ cells - This experiment is designed to  
determine whether mFLINT or anti-FasL antibody can improve  
15 the recovery of progenitor cells from purified human CD34+  
progenitor cells. Purified human CD34+ cells ( $1 \times 10^6$ /ml )  
are stimulated with 100 U/ml of human IL-6 and 100 ng/ml of  
human SCF, with or without mFLINT or anti-FasL antibody for  
3 days in vitro to expand progenitor cells. Following this  
20 expansion and treatment, cells are counted and a CFU assay  
is conducted.

#### EXAMPLE 26

##### TRANSGENIC FLINT MICE

##### 25 A - Preparation of Transgenic Mice

This example demonstrates the construction of  
transgenic mice expressing FLINT. These animals express  
significant levels of FLINT, yet demonstrate no adverse  
effects, which indicates that FLINT has a favorable toxicity  
30 profile. These animals are also useful in further

-91-

delineating useful treatment protocols for the conditions set out above.

*Transgene construction.* Polymerase chain reaction (PCR) primers were synthesized and used to amplify the fused  
5 FLINT-FLAG DNA sequences from plasmid pIG1-FLINTF, described above. The 5' primer, 5'-GAAGATCTTCTTTGATC

AAGGATGGGCTTCTGGACTT-3', a BglII restriction site and the 3' primer, 5'-GGACTAGTCCTGATCATCACTTGTCGTCGTCGTCCTT-3', contained an SpeI restriction enzyme site. These were used  
10 to amplify the entire FLINT and FLAG coding region. The amplified 1.1 kb fragment was ligated into the multiple cloning site of plasmid pMTmcs2 generating plasmid pMTmcs2-FLINT (6.7 kb). See Fox et al, Eur. J. Pharmacol. 308:195 (1996).

15 The FLINT gene fragment was excised from pMTmcs2-FLINT by BglII and SpeI digestion and gel-purified. This fragment then was blunted with Klenow enzyme and ligated into the Klenow-blunted MluI site of plasmid pLIV.7, provided by John Taylor of the J. David Gladstone Institutes. See Fan et  
20 al., Proc. Nat'l Acad. Sci. 91:8724 (1994). Resultant plasmid pLIV7-FLINT also contains the apo E gene promoter/5' flanking region and an hepatic enhancer sequences called the "hepatic control region" (HCR). For microinjection into embryos, a 7.0 kb DNA fragment encompassing the Apo E gene  
25 promoter-FLINT/FLAG-HCR fusion gene was excised from plasmid pLIV7-FLINT by digestion with SalI and SpeI and purified by gel electrophoresis and glass bead extraction.

*Transgenic animal development.* Transgenic mice were generated using established techniques described, for  
30 example, by Hogan, B. et al. (1986) MANIPULATING THE MOUSE EMBRYO: A LABORATORY MANUAL, Cold Spring Harbor Laboratory

-92-

(Cold Spring Harbor, NY), as modified by Fox and Solter, Molec. Cell. Biol. 8: 5470 (1988). Briefly, the 7.0 kb DNA fragment encompassing the Apo E gene promoter-FLINT-HCR fusion gene was microinjected into the male pronuclei of newly fertilized one-cell-stage embryos (zygotes) of the FVB/N strain. The embryos were cultured *in vitro* overnight to allow development to the two-cell-stage. Two-cell embryos were then transplanted into the oviducts of pseudopregnant CD-1 strain mice to allow development to term. To test for the presence of the transgene in the newborn mice, a small piece of toe was removed from each animal and digested with proteinase K to release the nucleic acids. A sample of the toe extract was subsequently subjected to PCR analysis using human FLINT-specific primers to identify transgene-containing mice. Five founder transgenic mice were identified as containing a FLINT transgene and were designated 6494, 7262, 7353, 7653 and 7659. Each of these founders was bred to produce stable lines of transgenics.

Line 6494 has been extensively characterized for transgene expression and pathology. Broad tissue expression was detected in line 6494 progeny by Northern blot, RT-PCR (TaqMan), Western blot and immunohistochemical analysis and was highest in the liver and kidney. High levels of FLINT protein were detected in the circulation of line 6494 mice by ELISA assay; founder level = 490 ng/ml; progeny levels ranged from 285-1360 ng/ml (n=6). No endogenous FLINT was detected by these analyses. No significant histopathologic findings were detected in 5 week or 8 week old 6494 progeny. Preliminary blood chemistry analysis suggested that triglyceride levels may be elevated in 6494 mice. The animals had no observable abnormalities.

-93-

B-Protection of mice from irradiation- This experiment is designed to evaluate whether transgenic FLINT mice are protected from the adverse effects of radiation. Lethally irradiate 18 transgenic and 18 non-transgenic mice with 850 cGy. Give no bone marrow transplant to 8 of the 18 mice per group. Of these 8 mice, use 5 as a control for radiation induced death and 3 for histological analysis of the intestinal mucosa and bone marrow compartments at 3 days post-irradiation.

Transplant the remaining 10 mice per group with  $3 \times 10^4$  bone marrow cells from normal non-transgenic donors. Since the liver is not damaged by irradiation, the FLINT transgenic mice should still be able to produce FLINT protein systemically. This bone marrow cell dose typically results in about 50% survival at 30 days for wild type animals that have been irradiated as described herein. Obtain 50-100  $\mu$ l of blood by tail bleed on days 7, 15, 21, and 30 days post-transplant and perform hematological analysis to monitor recovery of peripheral blood ( $n = 10$  mice per group; total of 20 mice). The hematological analysis will include a white blood cell count (WBC), red blood cell count (RBC), hematocrit, and blood smear microscopy to determine numbers of lymphocytes, neutrophils, eosinophils, platelets, mast cells and monocytes in the blood. Measure survival of mice and recovery of peripheral blood cells.

It is expected that, because FLINT prevents apoptosis of expanding progenitor cells and/or enhances recovery of intestinal epithelium from radiation-induced damage, there will be a difference between the two groups in survival,

-94-

histology of gut epithelium and bone marrow, and recovery of peripheral blood cells.

C-Protection of mice from chemotherapy- This experiment is designed to determine the effects of combining FLINT and GM-CSF administration on the recovery of progenitor cells from myelosuppression that is induced by sub-lethal irradiation and chemotherapy. In particular, this experiment examines whether white blood cell recovery, from a combination of chemotherapy and sublethal irradiation, is improved in transgenic FLINT mice. On example of such a chemotherapy regimen is treatment with carboplatin.

Administer 500 cGY of irradiation 4 hours following a single intraperitoneal injection of carboplatin (1.2 mg/mouse) to 15 transgenic and 15 control mice. This regimen induces severe myelosuppression with prolonged thrombocytopenia and severe anemia. Leonard et al., *Blood*, 83: 1499 (1994). Every day for 12 days, beginning 24 hours after irradiation, 10 µg/kg body weight of recombinant murine rmGM-CSF is injected i.p. to promote recovery of WBC count. Mayer et al., *J. Inf. Diseases* 163: 584 (1991), Gamba-Vitalo et al., 1991, *Blood Cells* 17: 193 (1991). Blood is collected by tail bleed every 7 days up to day 30 and a complete hematological analysis is conducted. The hematological analysis will include a white blood cell count (WBC), red blood cell count (RBC), hematocrit, and blood smear microscopy to determine numbers of lymphocytes, neutrophils, eosinophils, platelets, mast cells and monocytes in the blood. On days 12 and 20, 3 mice from each group are sacrificed. From these mice, spleen cellularity and bone marrow cellularity are analyzed. Also, the bone



-95-

marrow cells are pooled and used to set up CFU assays to examine progenitor content.

D- FLINT effects on peripheral mobilization of progenitor cells- This experiment is designed to test for testing peripheral mobilization of progenitor cells. To test for peripheral mobilization of progenitor cells, FLINT transgenic and control mice are treated with 3 mg of 5-FU i.p. and 4 days later, either CFU assays will be set up from isolated bone marrow cells, or 100ng of GM-CSF will be administered i.p. every day for 3 days followed by determination of bone marrow cellularity and progenitor content by CFU assays. The protocols for transplantation are described above in Example B. Applications for gene therapy would be to improve the recovery of progenitor cells following stimulation with cytokines either in vivo, or in vitro, for transduction with retrovirus, as described above.

#### EXAMPLE 27

##### Use of FLINT to Treat ALI Patient

A 55 year-old male presents to the emergency department unconscious. His family states that he was being treated as an outpatient for bronchitis for the past few days but worsened despite antibiotics. He has no relevant past history and his only medication was a third generation oral cephalosporin. Physical examination reveals an obtunded, cyanotic male who is hypotensive, tachypneic, and tachycardic, and who has bilateral lung congestion consistent with pulmonary edema. There is no evidence of congestive heart failure. Tests reveal hypoxemia (based on PaO<sub>2</sub>/FiO<sub>2</sub>) and bilateral lung infiltrates without cardiomegaly, consistent with a diagnosis of acute lung

-96-

injury. Based on the history it is concluded that the lung injury was a direct result of community-acquired pneumonia, and that the patient met the hypoxemia criteria for ALI within the last 12 hours. Ventilation measures include use of PEEP and low tidal volume. As soon as adequate oxygenation is confirmed, treatment with FLINT is initiated in the ER as an iv bolus of 2.5 mg/kg, followed by a continuous infusion of 0.1 mg/minute. FLINT along with aggressive supportive measures (e.g., positive ventilation, intravenous fluids, pressors, and nutritional support) are continued for four days in the ICU, at which time the FLINT is discontinued. Over the following 3 days, the patient begins to recover and is extubated on Day 8. He has an uneventful recovery and 6 months following discharge has no evidence of residual lung disease by blood gas and spirometry.

-97-

We claim:

1. mFLINT for use in treating acute respiratory distress syndrome.
- 5 2. mFLINT for use in treating ulcerative colitis.
3. mFLINT for use in inhibiting ulcerative colitis.
4. mFLINT for use in inhibiting ischemic injury during  
10 organ transplantation.
5. mFLINT for use in organ preservation during transplantation.
- 15 6. A formulation adapted for the treatment of acute respiratory distress syndrome, comprising mFLINT as an active ingredient.
7. A formulation adapted for the treatment of ulcerative  
20 colitis, comprising mFLINT as an active ingredient.
8. A formulation adapted for inhibiting ulcerative colitis, comprising mFLINT as an active ingredient.
- 25 9. A formulation adapted for inhibiting ischemic injury during organ transplantation comprising mFLINT as an active ingredient.
10. The use of mFLINT in the preparation of a medicament  
30 useful in treating acute respiratory distress syndrome.

-98-

11. The use of mFLINT in the preparation of a medicament useful in treating ulcerative colitis.

12. The use of mFLINT in the preparation of a medicament  
5 useful in inhibiting ulcerative colitis.

13. A method for treating acute respiratory distress syndrome which comprises administering a therapeutically effective amount of mFLINT to a mammal in need thereof.

10

14. A method for treating ulcerative colitis which comprises administering a therapeutically effective amount of mFLINT to a mammal in need thereof.

15 15. A method for inhibiting ulcerative colitis which comprises administering a therapeutically effective amount of mFLINT to a mammal in need thereof.

16. A liquid medium for infusion and preservation of organs  
20 said medium comprising mFLINT.

1/13

**FIG. 1a**

ATG AGG GCG CTG GAG GGG CCA GGC CTG TCG CTG TGC CTG GTG TTG Met Arg Ala Leu Glu Gly Pro Gly Leu Ser Leu Leu Cys Leu Val Leu	48
GCG CTG CCT GCC CTG CTG CCG GTG CCG GCT GTA CGC GGA GTG GCA GAA Ala Leu Pro Ala Leu Leu Pro Val Pro Ala Val Arg Gly Val Ala Glu	96
ACA CCC ACC TAC CCC TGG CGG GAC GCA GAG ACA GGG GAG CGG CTG GTG Thr Pro Thr Tyr Pro Trp Arg Asp Ala Glu Thr Gly Glu Arg Leu Val	144
TGC GCC CAG TGC CCC CCA GGC ACC TTT GTG CAG CGG CCG TGC CGC CGA Cys Ala Gln Cys Pro Pro Gly Thr Phe Val Gln Arg Pro Cys Arg Arg	192
GAC AGC CCC ACG ACG TGT GGC CCG TGT CCA CCG CGC CAC TAC ACG CAG Asp Ser Pro Thr Thr Cys Gly Pro Cys Pro Pro Arg His Tyr Thr Gln	240
TTC TGG AAC TAC CTG GAG CGC TGC CGC TAC TGC AAC GTC CTC TGC GGG Phe Trp Asn Tyr Leu Glu Arg Cys Arg Tyr Cys Asn Val Leu Cys Gly	288
GAG CGT GAG GAG GAG GCA CGG GCT TGC CAC GCC ACC CAC AAC CGT GCC Glu Arg Glu Glu Glu Ala Arg Ala Cys His Ala Thr His Asn Arg Ala 100 105 110	336
TGC CGC TGC CGC ACC GGC TTC TTC GCG CAC GCT GGT TTC TGC TTG GAG Cys Arg Cys Arg Thr Gly Phe Phe Ala His Ala Gly Phe Cys Leu Glu	384
CAC GCA TCG TGT CCA CCT GGT GCC GGC GTG ATT GCC CCG GGC ACC CCC His Ala Ser Cys Pro Pro Gly Ala Gly Val Ile Ala Pro Gly Thr Pro	432
AGC CAG AAC ACG CAG TGC CAG CCG TGC CCC CCA GGC ACC TTC TCA GCC Ser Gln Asn Thr Gln Cys Gln Pro Cys Pro Pro Gly Thr Phe Ser Ala	480
AGC AGC TCC AGC TCA GAG CAG TGC CAG CCC CAC CGC AAC TGC ACG GCC Ser Ser Ser Ser Ser Glu Gln Cys Gln Pro His Arg Asn Cys Thr Ala	528
CTG GGC CTG GCC CTC AAT GTG CCA GGC TCT TCC TCC CAT GAC ACC CTG Leu Gly Leu Ala Leu Asn Val Pro Gly Ser Ser Ser His Asp Thr Leu	576
TGC ACC AGC TGC ACT GGC TTC CCC CTC AGC ACC AGG GTA CCA GGA GCT Cys Thr Ser Cys Thr Gly Phe Pro Leu Ser Thr Arg Val Pro Gly Ala	624
GAG GAG TGT GAG CGT GCC GTC ATC GAC TTT GTG GCT TTC CAG GAC ATC Glu Glu Cys Glu Arg Ala Val Ile Asp Phe Val Ala Phe Gln Asp Ile	672
TCC ATC AAG AGG CTG CAG CGG CTG CTG CAG GCC CTC GAG GCC CCG GAG Ser Ile Lys Arg Leu Gln Arg Leu Leu Gln Ala Leu Glu Ala Pro Glu	720
GGC TGG GGT CCG ACA CCA AGG GCG GGC CGC GCG GCC TTG CAG CTG AAG Gly Trp Gly Pro Thr Pro Arg Ala Gly Arg Ala Ala Leu Gln Leu Lys	768

2/13

*FIG. 1b*

CTG CGT CGG CGG CTC ACG GAG CTC CTG GGG GCG CAG GAC GGG GCG CTG Leu Arg Arg Arg Leu Thr Glu Leu Leu Gly Ala Gln Asp Gly Ala Leu	816
CTG GTG CGG CTG CTG CAG GCG CTG CGC GTG GCC AGG ATG CCC GGG CTG Leu Val Arg Leu Leu Leu Gln Ala Leu Arg Val Ala Arg Met Pro Gly Leu	864
GAG CGG AGC GTC CGT GAG CGC TTC CTC CCT GTG CAC Glu Arg Ser Val Arg Glu Arg Phe Leu Pro Val His	900

3/13

**FIG. 2a**

gctctccctgctccagcaaggacc atg agg gcg ctg gag ggg cca ggc ctg	51
M R A L E G P G L	
tcg ctg ctg tgc ctg gtg ttg gcg ctg cct gcc ctg ctg ccg gtg	96
S L L C L V L A L P A L L P V	
ccg gct gta cgc gga gtg gca gaa aca ccc acc tac ccc tgg cgg	141
P A V R G V A E T P T Y P W R	
gac gca gag aca ggg gag cgg ctg gtg tgc gcc cag tgc ccc cca	186
D A E T G E R L V C A Q C P P	
ggc acc ttt gtg cag cgg ccg tgc cgc cga gac agc ccc acg acg	231
G T F V Q R P C R R D S P T T	
tgt ggc ccg tgt cca ccg cgc cac tac acg cag ttc tgg aac tac	276
C G P C P P R H Y T Q F W N Y	
ctg gag cgc tgc cgc tac tgc aac gtc ctc tgc ggg gag cgt gag	321
L E R C R Y C N V L C G E R E	
gag gag gca cgg gct tgc cac gcc acc cac aac cgt gcc tgc cgc	366
E E A R A C H A T H N R A C R	
tgc cgc acc ggc ttc ttc gcg cac gct ggt ttc tgc ttg gag cac	411
C R C R T G F F A H A G F C L	
gca tgc tgt cca cct ggt gcc ggc gtg att gcc ccg ggc acc ccc	456
E H A S C P P G A G V I A P G	
agc cag aac acg cag tgc cag ccg tgc ccc cca ggc acc ttc tca	501
T P S Q N T Q C Q P C P P G T	
gcc agc agc tcc agc tca gag cag tgc cag ccc cac cgc aac tgc	546
F S A S S S S S E Q C Q P H R	
acg gcc ctg ggc ctg gcc ctc aat gtg cca ggc tct tcc tcc cat	591
N C T A L G L A L N V P G S S	
gac acc ctg tgc acc agc tgc act ggc ttc ccc ctc agc acc agg	636
S H D T L C T S C T G F P L S	
gta cca gga gct gag gag tgt gag cgt gcc gtc atc gac ttt gtg	681
T R V F G A E E C E R A V I D	
gct ttc cag gac atc tcc atc aag agg ctg cag cgg ctg ctg cag	726
F V A F Q D I S I K R L Q R L	
gcc ctc gag gcc ccg gag ggc tgg gct ccg aca cca agg gcg ggc	771
L Q A L E A P E G W A P T P R	

4/13

*FIG. 2b*

cgc gcg gcc ttg cag ctg aag ctg cgt cgg cgg ctc acg gag ctc	816
A G R A A L Q L K L R R R L T	
ctg ggg gcg cag gac ggg gcg ctg ctg gtg cgg ctg ctg cag gcg	861
E L L G A Q D G A L L V R L L	
ctg cgc gtc gcc agg atg ccc ggg ctg gag cgg agc gtc cgt gag	906
Q A L R V A R M P G L E R S V	
cgc ctc ctc ccc gtc cac tga tcc tggccc	936
R E R F L P W H	



5/13

**FIG. 3**

gtg gca gaa aca ccc acc tac ccc tgg cgg gac gca gag aca ggg	45
V A E T P T Y P W R D A E T G	
gag cgg ctg gtg tgc gcc cag tgc ccc cca ggc acc ttt gtg cag	90
E R L V C A Q C P P G T F V Q	
cgg ccg tgc cgc cga gac agc ccc acg acg tgt ggc ccg tgt cca	135
R P C R R D S P T T C G P C P	
ccg cgc cac tac acg cag ttc tgg aac tac ctg gag cgc tgc cgc	180
P R H Y T Q F W N Y L E R C R	
tac tgc aac gtc ctc tgc ggg gag cgt gag gag gag gca cgg gct	225
Y C N V L C G E R E E E A R A	
tgc cac gcc acc cac aac cgt gcc tgc cgc tgc cgc acc ggc ttc	270
C H A T H N R A C R C R T G F	
ttc gcg cac gct ggt ttc tgc ttg gag cac gca tcg tgt cca cct	315
F A H A G F C L E H A S C P P	
ggt gcc ggc gtg att gcc ccg ggc acc ccc agc cag aac acg cag	360
G A G V I A P G T P S Q N T Q	
tgc cag ccg tgc ccc cca ggc acc ttc tca gcc agc agc tcc agc	405
C Q P C P P G T F S A S S S S	
tca gag cag tgc cag ccc cac cgc aac tgc acg gcc ctg ggc ctg	450
S E Q C Q P H R N C T A L G L	
gcc ctc aat gtg cca ggc tct tcc tcc cat gac acc ctg tgc acc	495
A L N V P G S S S H D T L C T	
agc tgc act ggc ttc ccc ctc agc acc agg gta cca gga gct gag	540
S C T G F P L S T R V P G A E	
gag tgt gag cgt gcc gtc atc gac ttt gtg gct ttc cag gac atc	585
E C E R A V I D F V A F Q D I	
tcc atc aag agg ctg cag cgg ctg ctg cag gcc ctc gag gcc ccg	630
S I K R L Q R L L Q A L E A P	
gag ggc tgg ggt ccg aca cca agg gcg ggc cgc gcg gcc ttg cag	675
E G W G P T P R A G R A A L Q	
ctg aag ctg cgt cgg cgg ctc acg gag ctc ctg ggg gcg cag gac	720
L K L R R R L T E L L G A Q D	
ggg gcg ctg ctg gtg cgg ctg ctg cag gcg ctg cgc gtg gcc agg	765
G A L L V R L L Q A L R V A R	
atg ccc ggg ctg gag cgg agc gtc cgt gag cgc ttc ctc cct gtg	810
M P G L E R S V R E R F L P V	
cac	813
H	

6/13

**FIG. 4a**

gtg gca gaa aca ccc acc tac ccc tgg cgg gac gca gag aca ggg	45
V A E T P T Y P W R D A E T G	
 gag cgg ctg gtg tgc gcc cag tgc ccc cca ggc acc ttt gtg cag	90
E R L V C A Q C P P G T F V Q	
 cgg ccg tgc cgc cga gac agc ccc acg acg tgt ggc ccg tgt cca	135
R P C R R D S P T T C G P C P	
 ccg cgc cac tac acg cag ttc tgg aac tac ctg gag cgc tgc cgc	180
P R H Y T Q F W N Y L E R C R	
 tac tgc aac gtc ctc tgc ggg gag cgt gag gag gag gca cgg gct	225
Y C N V L C G E R E E E A R A	
 tgc cac gcc acc cac aac cgt gcc tgc cgc tgc cgc acc ggc ttc	270
C H A T H N R A C R C R C R T	
 ttc gcg cac gct ggt ttc tgc ttg gag cac gca tcg tgt cca cct	315
G F F A H A G F C L E H A S C	
 ggt gcc ggc gtg att gcc ccg ggc acc ccc agc cag aac acg cag	360
P P G A G V I A P G T P S Q N	
 tgc cag ccg tgc ccc cca ggc acc ttc tca gcc agc agc tcc agc	405
T Q C Q P C P P G T F S A S S	
 tca gag cag tgc cag ccc cac cgc aac tgc acg gcc ctg ggc ctg	450
S S S E Q C Q P H R N C T A L	
 gcc ctc aat gtg cca ggc tct tcc tcc cat gac acc ctg tgc acc	495
G L A L N V P G S S S H D T L	
 agc tgc act ggc ttc ccc ctc agc acc agg gta cca gga gct gag	540
C T S C T G F P L S T R V P G	
 gag tgt gag cgt gcc gtc atc gac ttt gtg gct ttc cag gac atc	585
A E E C E R A V I D F V A F Q	
 tcc atc aag agg ctg cag cgg ctg ctg cag gcc ctc gag gcc ccg	630
D I S I K R L Q R L L Q A L E	
 gag ggc tgg gct ccg aca cca agg gcg ggc cgc gcg gcc ttg cag	675
A P E G W A P T P R A G R A A	
 ctg aag ctg cgt cgg cgg ctc acg gag ctc ctg ggg gcg cag gac	720
L Q L K L R R R L T E L L G A	

7/13

**FIG. 4b**

ggg gcg ctg ctg gtc cgg ctg cag gcg ctg cgc gtc gcc agg	765
Q D G A L L V R L L Q A L R V	
atg ccc ggg ctg gag cgg agc gtc cgt gag cgc ttc ctc cct gtc	810
A R M P G L E R S V R E R F L	
cac tga tcc tggccc	825
P W H	

8/13

FIG. 5a





FIG. 5b



9/13

FIG. 6a




FIG. 6b






FIG. 6c




10/13


**FIG. 7**



II/13

**FIG. 8**


**FIG. 9**  
FLINT IMPROVES RECOVERY OF BM PROGENITOR CELLS  
DURING CYTOKINE STIMULATION IN VITRO





13/13

**FIG. 10**  
FLINT OR ANTI-FasL IMPROVES RECOVERY OF BONE  
MARROW CELLS FOLLOWING GROWTH FACTOR STIMULATION



## SEQUENCE LISTING

&lt;110&gt; ELI LILLY AND COMPANY

&lt;120&gt; THERAPEUTIC APPLICATIONS OF FLINT POLYPEPTIDES

&lt;130&gt; X-12915A

&lt;150&gt; US 09/280,567

&lt;151&gt; 1999-03-30

&lt;160&gt; 13

&lt;170&gt; PatentIn Ver. 2.0

&lt;210&gt; 1

&lt;211&gt; 900

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)...(900)

&lt;400&gt; 1

atg	agg	gcg	ctg	gag	ggg	cca	ggc	ctg	tcg	ctg	ctg	tgc	ctg	gtg	ttg	48
Met	Arg	Ala	Leu	Glu	Gly	Pro	Gly	Leu	Ser	Leu	Leu	Cys	Leu	Val	Leu	
1				5				10					15			

gcg	ctg	cct	gcc	ctg	ctg	ccg	gtg	ccg	gct	gta	cgc	gga	gtg	gca	gaa	96
Ala	Leu	Pro	Ala	Leu	Leu	Pro	Val	Pro	Ala	Val	Arg	Gly	Val	Ala	Glu	
			20				25						30			

aca	ccc	acc	tac	ccc	tgg	cgg	gac	gca	gag	aca	ggg	gag	cgg	ctg	gtg	144
Thr	Pro	Thr	Tyr	Pro	Trp	Arg	Asp	Ala	Glu	Thr	Gly	Glu	Arg	Leu	Val	
		35				40						45				

tgc	gcc	cag	tgc	ccc	cca	ggc	acc	ttt	gtg	cag	cgg	ccg	tgc	cgc	cga	192
Cys	Ala	Gln	Cys	Pro	Pro	Gly	Thr	Phe	Val	Gln	Arg	Pro	Cys	Arg	Arg	
	50					55					60					

gac	agc	ccc	acg	acg	tgt	ggc	ccg	tgt	cca	ccg	cgc	cac	tac	acg	cag	240
Asp	Ser	Pro	Thr	Thr	Cys	Gly	Pro	Cys	Pro	Pro	Arg	His	Tyr	Thr	Gln	
	65				70				75					80		

ttc	tgg	aac	tac	ctg	gag	cgc	tgc	cgc	tac	tgc	aac	gtc	ctc	tgc	ggg	288
Phe	Trp	Asn	Tyr	Leu	Glu	Arg	Cys	Arg	Tyr	Cys	Asn	Val	Leu	Cys	Gly	
			85					90						95		

gag	cgt	gag	gag	gag	gca	cgg	gct	tgc	cac	gcc	acc	cac	aac	cgt	gcc	336
Glu	Arg	Glu	Glu	Glu	Ala	Arg	Ala	Cys	His	Ala	Thr	His	Asn	Arg	Ala	
			100				105						110			

tgc	cgc	tgc	cgc	acc	ggc	ttc	ttc	gcg	cac	gct	ggt	ttc	tgc	ttg	gag	384
Cys	Arg	Cys	Arg	Thr	Gly	Phe	Phe	Ala	His	Ala	Gly	Phe	Cys	Leu	Glu	
			115			120						125				

```

cac gca tgc tgt cca cct ggt gcc ggc gtg att gcc ccg ggc acc ccc 432
His Ala Ser Cys Pro Pro Gly Ala Gly Val Ile Ala Pro Gly Thr Pro
130 135 140

agc cag aac acg cag tgc cag ccg tgc ccc cca ggc acc ttc tca gcc 480
Ser Gln Asn Thr Gln Cys Gln Pro Cys Pro Pro Gly Thr Phe Ser Ala
145 150 155 160

agc agc tcc agc tca gag cag tgc cag ccc cac cgc aac tgc acg gcc 528
Ser Ser Ser Ser Ser Glu Gln Cys Gln Pro His Arg Asn Cys Thr Ala
165 170 175

ctg ggc ctg gcc ctc aat gtg cca ggc tct tcc tcc cat gac acc ctg 576
Leu Gly Leu Ala Leu Asn Val Pro Gly Ser Ser Ser His Asp Thr Leu
180 185 190

tgc acc agc tgc act ggc ttc ccc ctc agc acc agg gta cca gga gct 624
Cys Thr Ser Cys Thr Gly Phe Pro Leu Ser Thr Arg Val Pro Gly Ala
195 200 205

gag gag tgt gag cgt gcc gtc atc gac ttt gtg gct ttc cag gac atc 672
Glu Glu Cys Glu Arg Ala Val Ile Asp Phe Val Ala Phe Gln Asp Ile
210 215 220

tcc atc aag agg ctg cag cgg ctg ctg cag gcc ctc gag gcc ccg gag 720
Ser Ile Lys Arg Leu Gln Arg Leu Leu Gln Ala Leu Glu Ala Pro Glu
225 230 235 240

ggc tgg ggt ccg aca cca agg gcg ggc cgc gcg gcc ttg cag ctg aag 768
Gly Trp Gly Pro Thr Pro Arg Ala Gly Arg Ala Ala Leu Gln Leu Lys
245 250 255

ctg cgt cgg cgg ctc acg gag ctc ctg ggg gcg cag gac ggg gcg ctg 816
Leu Arg Arg Arg Leu Thr Glu Leu Leu Gly Ala Gln Asp Gly Ala Leu
260 265 270

ctg gtg cgg ctg ctg cag gcg ctg cgc gtg gcc agg atg ccc ggg ctg 864
Leu Val Arg Leu Leu Gln Ala Leu Arg Val Ala Arg Met Pro Gly Leu
275 280 285

gag cgg agc gtc cgt gag cgc ttc ctc cct gtg cac 900
Glu Arg Ser Val Arg Glu Arg Phe Leu Pro Val His
290 295 300

```

&lt;210&gt; 2

&lt;211&gt; 300

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 2

```

Met Arg Ala Leu Glu Gly Pro Gly Leu Ser Leu Leu Cys Leu Val Leu
1 5 10 15

```

```

Ala Leu Pro Ala Leu Leu Pro Val Pro Ala Val Arg Gly Val Ala Glu
20 25 30

```

Thr Pro Thr Tyr Pro Trp Arg Asp Ala Glu Thr Gly Glu Arg Leu Val  
                   35                                  40                                  45  
 Cys Ala Gln Cys Pro Pro Gly Thr Phe Val Gln Arg Pro Cys Arg Arg  
                   50                                  55                                  60  
 Asp Ser Pro Thr Thr Cys Gly Pro Cys Pro Pro Arg His Tyr Thr Gln  
                   65                                  70                                  75                                  80  
 Phe Trp Asn Tyr Leu Glu Arg Cys Arg Tyr Cys Asn Val Leu Cys Gly  
                                   85                                  90                                  95  
 Glu Arg Glu Glu Glu Ala Arg Ala Cys His Ala Thr His Asn Arg Ala  
                                   100                                  105                                  110  
 Cys Arg Cys Arg Thr Gly Phe Phe Ala His Ala Gly Phe Cys Leu Glu  
                                   115                                  120                                  125  
 His Ala Ser Cys Pro Pro Gly Ala Gly Val Ile Ala Pro Gly Thr Pro  
                                   130                                  135                                  140  
 Ser Gln Asn Thr Gln Cys Gln Pro Cys Pro Pro Gly Thr Phe Ser Ala  
                                   145                                  150                                  155                                  160  
 Ser Ser Ser Ser Ser Glu Gln Cys Gln Pro His Arg Asn Cys Thr Ala  
                                   165                                  170                                  175  
 Leu Gly Leu Ala Leu Asn Val Pro Gly Ser Ser Ser His Asp Thr Leu  
                                   180                                  185                                  190  
 Cys Thr Ser Cys Thr Gly Phe Pro Leu Ser Thr Arg Val Pro Gly Ala  
                                   195                                  200                                  205  
 Glu Glu Cys Glu Arg Ala Val Ile Asp Phe Val Ala Phe Gln Asp Ile  
                                   210                                  215                                  220  
 Ser Ile Lys Arg Leu Gln Arg Leu Leu Gln Ala Leu Glu Ala Pro Glu  
                                   225                                  230                                  235                                  240  
 Gly Trp Gly Pro Thr Pro Arg Ala Gly Arg Ala Ala Leu Gln Leu Lys  
                                   245                                  250                                  255  
 Leu Arg Arg Arg Leu Thr Glu Leu Leu Gly Ala Gln Asp Gly Ala Leu  
                                   260                                  265                                  270  
 Leu Val Arg Leu Leu Gln Ala Leu Arg Val Ala Arg Met Pro Gly Leu  
                                   275                                  280                                  285  
 Glu Arg Ser Val Arg Glu Arg Phe Leu Pro Val His  
                                   290                                  295                                  300

&lt;210&gt; 3

&lt;211&gt; 936

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (25)..(924)

&lt;400&gt; 3

```

gctctccctg ctccagcaag gacc atg agg gcg ctg gag ggg cca ggc ctg 51
      Met Arg Ala Leu Glu Gly Pro Gly Leu
              1              5

tcg ctg ctg tgc ctg gtg ttg gcg ctg cct gcc ctg ctg ccg gtg ccg 99
Ser Leu Leu Cys Leu Val Leu Ala Leu Pro Ala Leu Leu Pro Val Pro
10              15              20              25

gct gta cgc gga gtg gca gaa aca ccc acc tac ccc tgg cgg gac gca 147
Ala Val Arg Gly Val Ala Glu Thr Pro Thr Tyr Pro Trp Arg Asp Ala
30              35              40

gag aca ggg gag cgg ctg gtg tgc gcc cag tgc ccc cca ggc acc ttt 195
Glu Thr Gly Glu Arg Leu Val Cys Ala Gln Cys Pro Pro Gly Thr Phe
45              50              55

gtg cag cgg ccg tgc cgc cga gac agc ccc acg acg tgt ggc ccg tgt 243
Val Gln Arg Pro Cys Arg Arg Asp Ser Pro Thr Thr Cys Gly Pro Cys
60              65              70

cca ccg cgc cac tac acg cag ttc tgg aac tac ctg gag cgc tgc cgc 291
Pro Pro Arg His Tyr Thr Gln Phe Trp Asn Tyr Leu Glu Arg Cys Arg
75              80              85

tac tgc aac gtc ctc tgc ggg gag cgt gag gag gag gca cgg gct tgc 339
Tyr Cys Asn Val Leu Cys Gly Glu Arg Glu Glu Glu Ala Arg Ala Cys
90              95              100              105

cac gcc acc cac aac cgt gcc tgc cgc tgc cgc acc ggc ttc ttc gcg 387
His Ala Thr His Asn Arg Ala Cys Arg Cys Arg Thr Gly Phe Phe Ala
110              115              120

cac gct ggt ttc tgc ttg gag cac gca tcg tgt cca cct ggt gcc ggc 435
His Ala Gly Phe Cys Leu Glu His Ala Ser Cys Pro Pro Gly Ala Gly
125              130              135

gtg att gcc ccg ggc acc ccc agc cag aac acg cag tgc cag ccg tgc 483
Val Ile Ala Pro Gly Thr Pro Ser Gln Asn Thr Gln Cys Gln Pro Cys
140              145              150

ccc cca ggc acc ttc tca gcc agc agc tcc agc tca gag cag tgc cag 531
Pro Pro Gly Thr Phe Ser Ala Ser Ser Ser Ser Ser Glu Gln Cys Gln
155              160              165

ccc cac cgc aac tgc acg gcc ctg ggc ctg gcc ctc att gtg cca ggc 579
Pro His Arg Asn Cys Thr Ala Leu Gly Leu Ala Leu Ile Val Pro Gly
170              175              180              185

tct tcc tcc cat gac acc ctg tgc acc agc tgc act ggc ttc ccc ctc 627
Ser Ser Ser His Asp Thr Leu Cys Thr Ser Cys Thr Gly Phe Pro Leu
190              195              200

```

agc acc agg gta cca gga gct gag gag tgt gag cgt gcc gtc atc gac 675  
 Ser Thr Arg Val Pro Gly Ala Glu Glu Cys Glu Arg Ala Val Ile Asp  
 205 210 215

ttt gtg gct ttc cag gac atc tcc atc aag agg ctg cag cgg ctg ctg 723  
 Phe Val Ala Phe Gln Asp Ile Ser Ile Lys Arg Leu Gln Arg Leu Leu  
 220 225 230

cag gcc ctc gag gcc ccg gag ggc tgg gct ccg aca cca agg gcg ggc 771  
 Gln Ala Leu Glu Ala Pro Glu Gly Trp Ala Pro Thr Pro Arg Ala Gly  
 235 240 245

cgc gcg gcc ttg cag ctg aag ctg cgt cgg cgg ctc acg gag ctc ctg 819  
 Arg Ala Ala Leu Gln Leu Lys Leu Arg Arg Arg Leu Thr Glu Leu Leu  
 250 255 260 265

ggg gcg cag gac ggg gcg ctg ctg gtg cgg ctg ctg cag gcg ctg cgc 867  
 Gly Ala Gln Asp Gly Ala Leu Leu Val Arg Leu Leu Gln Ala Leu Arg  
 270 275 280

gtg gcc agg atg ccc ggg ctg gag cgg agc gtc cgt gag cgc ttc ctc 915  
 Val Ala Arg Met Pro Gly Leu Glu Arg Ser Val Arg Glu Arg Phe Leu  
 285 290 295

cct gtg cac tgatcctggc cc 936  
 Pro Val His  
 300

<210> 4  
 <211> 300  
 <212> PRT  
 <213> Homo sapiens

<400> 4  
 Met Arg Ala Leu Glu Gly Pro Gly Leu Ser Leu Leu Cys Leu Val Leu  
 1 5 10 15

Ala Leu Pro Ala Leu Leu Pro Val Pro Ala Val Arg Gly Val Ala Glu  
 20 25 30

Thr Pro Thr Tyr Pro Trp Arg Asp Ala Glu Thr Gly Glu Arg Leu Val  
 35 40 45

Cys Ala Gln Cys Pro Pro Gly Thr Phe Val Gln Arg Pro Cys Arg Arg  
 50 55 60

Asp Ser Pro Thr Thr Cys Gly Pro Cys Pro Pro Arg His Tyr Thr Gln  
 65 70 75 80

Phe Trp Asn Tyr Leu Glu Arg Cys Arg Tyr Cys Asn Val Leu Cys Gly  
 85 90 95

Glu Arg Glu Glu Glu Ala Arg Ala Cys His Ala Thr His Asn Arg Ala  
 100 105 110

Cys Arg Cys Arg Thr Gly Phe Phe Ala His Ala Gly Phe Cys Leu Glu

115                      120                      125  
 His Ala Ser Cys Pro Pro Gly Ala Gly Val Ile Ala Pro Gly Thr Pro  
 130                      135                      140  
 Ser Gln Asn Thr Gln Cys Gln Pro Cys Pro Pro Gly Thr Phe Ser Ala  
 145                      150                      155                      160  
 Ser Ser Ser Ser Ser Glu Gln Cys Gln Pro His Arg Asn Cys Thr Ala  
 165                      170                      175  
 Leu Gly Leu Ala Leu Ile Val Pro Gly Ser Ser Ser His Asp Thr Leu  
 180                      185                      190  
 Cys Thr Ser Cys Thr Gly Phe Pro Leu Ser Thr Arg Val Pro Gly Ala  
 195                      200                      205  
 Glu Glu Cys Glu Arg Ala Val Ile Asp Phe Val Ala Phe Gln Asp Ile  
 210                      215                      220  
 Ser Ile Lys Arg Leu Gln Arg Leu Leu Gln Ala Leu Glu Ala Pro Glu  
 225                      230                      235                      240  
 Gly Trp Ala Pro Thr Pro Arg Ala Gly Arg Ala Ala Leu Gln Leu Lys  
 245                      250                      255  
 Leu Arg Arg Arg Leu Thr Glu Leu Leu Gly Ala Gln Asp Gly Ala Leu  
 260                      265                      270  
 Leu Val Arg Leu Leu Gln Ala Leu Arg Val Ala Arg Met Pro Gly Leu  
 275                      280                      285  
 Glu Arg Ser Val Arg Glu Arg Phe Leu Pro Val His  
 290                      295                      300

<210> 5  
 <211> 813  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (1)..(813)

<400> 5  
 gtg gca gaa aca ccc acc tac ccc tgg cgg gac gca gag aca ggg gag 48  
 Val Ala Glu Thr Pro Thr Tyr Pro Trp Arg Asp Ala Glu Thr Gly Glu  
 1                      5                      10                      15  
 cgg ctg gtg tgc gcc cag tgc ccc cca ggc acc ttt gtg cag cgg ccg 96  
 Arg Leu Val Cys Ala Gln Cys Pro Pro Gly Thr Phe Val Gln Arg Pro  
 20                      25                      30  
 tgc cgc cga gac agc ccc acg acg tgt ggc ccg tgt cca ccg cgc cac 144  
 Cys Arg Arg Asp Ser Pro Thr Thr Cys Gly Pro Cys Pro Pro Arg His  
 35                      40                      45

tac acg cag ttc tgg aac tac ctg gag cgc tgc cgc tac tgc aac gtc	192
Tyr Thr Gln Phe Trp Asn Tyr Leu Glu Arg Cys Arg Tyr Cys Asn Val	
50 55 60	
ctc tgc ggg gag cgt gag gag gag gca cgg gct tgc cac gcc acc cac	240
Leu Cys Gly Glu Arg Glu Glu Glu Ala Arg Ala Cys His Ala Thr His	
65 70 75 80	
aac cgt gcc tgc cgc tgc cgc acc ggc ttc ttc gcg cac gct ggt ttc	288
Asn Arg Ala Cys Arg Cys Arg Thr Gly Phe Phe Ala His Ala Gly Phe	
85 90 95	
tgc ttg gag cac gca tgc tgt cca cct ggt gcc ggc gtg att gcc ccg	336
Cys Leu Glu His Ala Ser Cys Pro Pro Gly Ala Gly Val Ile Ala Pro	
100 105 110	
ggc acc ccc agc cag aac acg cag tgc cag ccg tgc ccc cca ggc acc	384
Gly Thr Pro Ser Gln Asn Thr Gln Cys Gln Pro Cys Pro Pro Gly Thr	
115 120 125	
ttc tca gcc agc agc tcc agc tca gag cag tgc cag ccc cac cgc aac	432
Phe Ser Ala Ser Ser Ser Ser Ser Glu Gln Cys Gln Pro His Arg Asn	
130 135 140	
tgc acg gcc ctg ggc ctg gcc ctc aat gtg cca ggc tct tcc tcc cat	480
Cys Thr Ala Leu Gly Leu Ala Leu Asn Val Pro Gly Ser Ser Ser His	
145 150 155 160	
gac acc ctg tgc acc agc tgc act ggc ttc ccc ctc agc acc agg gta	528
Asp Thr Leu Cys Thr Ser Cys Thr Gly Phe Pro Leu Ser Thr Arg Val	
165 170 175	
cca gga gct gag gag tgt gag cgt gcc gtc atc gac ttt gtg gct ttc	576
Pro Gly Ala Glu Glu Cys Glu Arg Ala Val Ile Asp Phe Val Ala Phe	
180 185 190	
cag gac atc tcc atc aag agg ctg cag cgg ctg ctg cag gcc ctc gag	624
Gln Asp Ile Ser Ile Lys Arg Leu Gln Arg Leu Leu Gln Ala Leu Glu	
195 200 205	
gcc ccg gag ggc tgg ggt ccg aca cca agg gcg ggc cgc gcg gcc ttg	672
Ala Pro Glu Gly Trp Gly Pro Thr Pro Arg Ala Gly Arg Ala Ala Leu	
210 215 220	
cag ctg aag ctg cgt cgg cgg ctc acg gag ctc ctg ggg gcg cag gac	720
Gln Leu Lys Leu Arg Arg Arg Leu Thr Glu Leu Leu Gly Ala Gln Asp	
225 230 235 240	
ggg gcg ctg ctg gtg cgg ctg ctg cag gcg ctg cgc gtg gcc agg atg	768
Gly Ala Leu Leu Val Arg Leu Leu Gln Ala Leu Arg Val Ala Arg Met	
245 250 255	
ccc ggg ctg gag cgg agc gtc cgt gag cgc ttc ctc cct gtg cac	813
Pro Gly Leu Glu Arg Ser Val Arg Glu Arg Phe Leu Pro Val His	
260 265 270	



<210> 6  
 <211> 271  
 <212> PRT  
 <213> Homo sapiens

<400> 6

```

Val Ala Glu Thr Pro Thr Tyr Pro Trp Arg Asp Ala Glu Thr Gly Glu
  1                      5                      10                      15

Arg Leu Val Cys Ala Gln Cys Pro Pro Gly Thr Phe Val Gln Arg Pro
      20                      25                      30

Cys Arg Arg Asp Ser Pro Thr Thr Cys Gly Pro Cys Pro Pro Arg His
      35                      40                      45

Tyr Thr Gln Phe Trp Asn Tyr Leu Glu Arg Cys Arg Tyr Cys Asn Val
      50                      55                      60

Leu Cys Gly Glu Arg Glu Glu Glu Ala Arg Ala Cys His Ala Thr His
      65                      70                      75                      80

Asn Arg Ala Cys Arg Cys Arg Thr Gly Phe Phe Ala His Ala Gly Phe
      85                      90                      95

Cys Leu Glu His Ala Ser Cys Pro Pro Gly Ala Gly Val Ile Ala Pro
      100                      105                      110

Gly Thr Pro Ser Gln Asn Thr Gln Cys Gln Pro Cys Pro Pro Gly Thr
      115                      120                      125

Phe Ser Ala Ser Ser Ser Ser Ser Glu Gln Cys Gln Pro His Arg Asn
      130                      135                      140

Cys Thr Ala Leu Gly Leu Ala Leu Asn Val Pro Gly Ser Ser Ser His
      145                      150                      155                      160

Asp Thr Leu Cys Thr Ser Cys Thr Gly Phe Pro Leu Ser Thr Arg Val
      165                      170                      175

Pro Gly Ala Glu Glu Cys Glu Arg Ala Val Ile Asp Phe Val Ala Phe
      180                      185                      190

Gln Asp Ile Ser Ile Lys Arg Leu Gln Arg Leu Leu Gln Ala Leu Glu
      195                      200                      205

Ala Pro Glu Gly Trp Gly Pro Thr Pro Arg Ala Gly Arg Ala Ala Leu
      210                      215                      220

Gln Leu Lys Leu Arg Arg Arg Leu Thr Glu Leu Leu Gly Ala Gln Asp
      225                      230                      235                      240

Gly Ala Leu Leu Val Arg Leu Leu Gln Ala Leu Arg Val Ala Arg Met
      245                      250                      255

Pro Gly Leu Glu Arg Ser Val Arg Glu Arg Phe Leu Pro Val His
      260                      265                      270

```

<210> 7  
 <211> 825  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (1)..(813)

```

<400> 7
gtg gca gaa aca ccc acc tac ccc tgg cgg gac gca gag aca ggg gag      48
Val Ala Glu Thr Pro Thr Tyr Pro Trp Arg Asp Ala Glu Thr Gly Glu
   1                      5                      10                      15

cgg ctg gtg tgc gcc cag tgc ccc cca ggc acc ttt gtg cag cgg ccg      96
Arg Leu Val Cys Ala Gln Cys Pro Pro Gly Thr Phe Val Gln Arg Pro
                20                      25                      30

tgc cgc cga gac agc ccc acg acg tgt ggc ccg tgt cca ccg cgc cac      144
Cys Arg Arg Asp Ser Pro Thr Thr Cys Gly Pro Cys Pro Pro Arg His
                35                      40                      45

tac acg cag ttc tgg aac tac ctg gag cgc tgc cgc tac tgc aac gtc      192
Tyr Thr Gln Phe Trp Asn Tyr Leu Glu Arg Cys Arg Tyr Cys Asn Val
   50                      55                      60

ctc tgc ggg gag cgt gag gag gag gca cgg gct tgc cac gcc acc cac      240
Leu Cys Gly Glu Arg Glu Glu Glu Ala Arg Ala Cys His Ala Thr His
   65                      70                      75                      80

aac cgt gcc tgc cgc tgc cgc acc ggc ttc ttc gcg cac gct ggt ttc      288
Asn Arg Ala Cys Arg Cys Arg Thr Gly Phe Phe Ala His Ala Gly Phe
                85                      90                      95

tgc ttg gag cac gca tcg tgt cca cct ggt gcc ggc gtg att gcc ccg      336
Cys Leu Glu His Ala Ser Cys Pro Pro Gly Ala Gly Val Ile Ala Pro
                100                      105                      110

ggc acc ccc agc cag aac acg cag tgc cag ccg tgc ccc cca ggc acc      384
Gly Thr Pro Ser Gln Asn Thr Gln Cys Gln Pro Cys Pro Pro Gly Thr
                115                      120                      125

ttc tca gcc agc agc tcc agc tca gag cag tgc cag ccc cac cgc aac      432
Phe Ser Ala Ser Ser Ser Ser Ser Glu Gln Cys Gln Pro His Arg Asn
                130                      135                      140

tgc acg gcc ctg ggc ctg gcc ctc aat gtg cca ggc tct tcc tcc cat      480
Cys Thr Ala Leu Gly Leu Ala Leu Asn Val Pro Gly Ser Ser Ser His
   145                      150                      155                      160

gac acc ctg tgc acc agc tgc act ggc ttc ccc ctc agc acc agg gta      528
Asp Thr Leu Cys Thr Ser Cys Thr Gly Phe Pro Leu Ser Thr Arg Val
                165                      170                      175

cca gga gct gag gag tgt gag cgt gcc gtc atc gac ttt gtg gct ttc      576

```

Pro Gly Ala Glu Glu Cys Glu Arg Ala Val Ile Asp Phe Val Ala Phe  
 180 185 190

cag gac atc tcc atc aag agg ctg cag cgg ctg ctg cag gcc ctc gag 624  
 Gln Asp Ile Ser Ile Lys Arg Leu Gln Arg Leu Leu Gln Ala Leu Glu  
 195 200 205

gcc ccg gag ggc tgg gct ccg aca cca agg gcg ggc cgc gcg gcc ttg 672  
 Ala Pro Glu Gly Trp Ala Pro Thr Pro Arg Ala Gly Arg Ala Ala Leu  
 210 215 220

cag ctg aag ctg cgt cgg cgg ctc acg gag ctc ctg ggg gcg cag gac 720  
 Gln Leu Lys Leu Arg Arg Arg Leu Thr Glu Leu Leu Gly Ala Gln Asp  
 225 230 235 240

ggg gcg ctg ctg gtg cgg ctg ctg cag gcg ctg cgc gtg gcc agg atg 768  
 Gly Ala Leu Leu Val Arg Leu Leu Gln Ala Leu Arg Val Ala Arg Met  
 245 250 255

ccc ggg ctg gag cgg agc gtc cgt gag cgc ttc ctc cct gtg cac 813  
 Pro Gly Leu Glu Arg Ser Val Arg Glu Arg Phe Leu Pro Val His  
 260 265 270

tgatcctggc cc 825

<210> 8  
 <211> 271  
 <212> PRT  
 <213> Homo sapiens

<400> 8  
 Val Ala Glu Thr Pro Thr Tyr Pro Trp Arg Asp Ala Glu Thr Gly Glu  
 1 5 10 15

Arg Leu Val Cys Ala Gln Cys Pro Pro Gly Thr Phe Val Gln Arg Pro  
 20 25 30

Cys Arg Arg Asp Ser Pro Thr Thr Cys Gly Pro Cys Pro Pro Arg His  
 35 40 45

Tyr Thr Gln Phe Trp Asn Tyr Leu Glu Arg Cys Arg Tyr Cys Asn Val  
 50 55 60

Leu Cys Gly Glu Arg Glu Glu Glu Ala Arg Ala Cys His Ala Thr His  
 65 70 75 80

Asn Arg Ala Cys Arg Cys Arg Thr Gly Phe Phe Ala His Ala Gly Phe  
 85 90 95

Cys Leu Glu His Ala Ser Cys Pro Pro Gly Ala Gly Val Ile Ala Pro  
 100 105 110

Gly Thr Pro Ser Gln Asn Thr Gln Cys Gln Pro Cys Pro Pro Gly Thr  
 115 120 125

Phe Ser Ala Ser Ser Ser Ser Ser Glu Gln Cys Gln Pro His Arg Asn

130                      135                      140  
 Cys Thr Ala Leu Gly Leu Ala Leu Asn Val Pro Gly Ser Ser Ser His  
 145                      150                      155                      160  
 Asp Thr Leu Cys Thr Ser Cys Thr Gly Phe Pro Leu Ser Thr Arg Val  
                     165                      170                      175  
 Pro Gly Ala Glu Glu Cys Glu Arg Ala Val Ile Asp Phe Val Ala Phe  
                     180                      185                      190  
 Gln Asp Ile Ser Ile Lys Arg Leu Gln Arg Leu Leu Gln Ala Leu Glu  
                     195                      200                      205  
 Ala Pro Glu Gly Trp Ala Pro Thr Pro Arg Ala Gly Arg Ala Ala Leu  
                     210                      215                      220  
 Gln Leu Lys Leu Arg Arg Arg Leu Thr Glu Leu Leu Gly Ala Gln Asp  
 225                      230                      235                      240  
 Gly Ala Leu Leu Val Arg Leu Leu Gln Ala Leu Arg Val Ala Arg Met  
                     245                      250                      255  
 Pro Gly Leu Glu Arg Ser Val Arg Glu Arg Phe Leu Pro Val His  
                     260                      265                      270

<210> 9  
 <211> 8  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Polypeptide

<400> 9  
 Asp Tyr Lys Asp Asp Asp Asp Lys  
   1                                  5

<210> 10  
 <211> 59  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 10  
 tagggctgat caaggatggg cttctggact tgggcggccc ctccgcaggc ggaccgggg 59

<210> 11  
 <211> 66  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 11

aggggggcg cgcgtgatca tcacttgctg tcgtcgtcct tgtagtcgtg cacagggagg 60

aagcgc

66

<210> 12

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 12

gaagatcttc ttgatcaag gatgggcttc tggactt

37

<210> 13

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 13

ggactagtcc tgatcatcac ttgtcgtcgt cgtcctt

37

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>A61K 38/17</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 00/37094</b> <b>(43) International Publication Date:</b> 29 June 2000 (29.06.00)
<b>(21) International Application Number:</b> PCT/US99/30734 <b>(22) International Filing Date:</b> 21 December 1999 (21.12.99)  <b>(30) Priority Data:</b> 60/113,407 22 December 1998 (22.12.98) US PCT/US99/06797 30 March 1999 (30.03.99) US 60/172,239 20 October 1999 (20.10.99) US  <b>(71) Applicant (for all designated States except US):</b> ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> COHEN, Fredric, Jay [US/US]; 8330 Catamaran Drive, Indianapolis, IN 46236 (US). POSADA, James, Arthur [US/US]; 3575 Inverness Boulevard, Carmel, IN 46032 (US). WIERDA, Daniel [US/US]; 2606 South Brandywine Trail, Greenfield, IN 46140 (US).  <b>(74) Agents:</b> WEBSTER, Thomas, D. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>  <b>(88) Date of publication of the international search report:</b> 9 November 2000 (09.11.00)
<b>(54) Title:</b> THERAPEUTIC APPLICATIONS OF FLINT POLYPEPTIDES  <b>(57) Abstract</b>  Mature FLINT protein (mFLINT) binds FasL and LIGHT, and prevents FasL-Fas interaction. mFLINT inhibits FasL-Fas-mediated apoptotic and proinflammatory activity, and is useful in treating disorders that may be associated with abnormal apoptosis and inflammation. The invention provides the amino acid and nucleotide sequences of FLINT and mature FLINT. Therapeutic compositions and methods of treatment utilizing mFLINT also are provided.		

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/30734

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 A61K38/17

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 861 850 A (SMITHKLINE BEECHAM CORP) 2 September 1998 (1998-09-02) page 8, line 36 - line 40 page 10, line 30 - line 51; claims Sequence ID no.2 page 11; example 1 ---	1,6,10, 13
X	WO 98 30694 A (HUMAN GENOME SCIENCES INC ;FENG PING (US); NI JIAN (US); EBNER REI) 16 July 1998 (1998-07-16) page 39, line 5 -page 48, line 2; claims ---	1
A	---	6,10,13
P,X	WO 99 50413 A (MIZRAHI JACQUES ;HEUER JOSEF GEORG (US); NOBLITT TIMOTHY WAYNE (US) 7 October 1999 (1999-10-07) claims; example 11 ---	1,6,10, 13
	-/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents :

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*Z\* document member of the same patent family

Date of the actual completion of the international search

11 May 2000

Date of mailing of the international search report

24.07.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

LE CORNEC N.D.R.



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/30734

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 99 14330 A (GENENTECH INC) 25 March 1999 (1999-03-25)	1
P,A	page 25, line 22 -page 27, line 44; claims ---	6,10,13
P,X	WO 99 07738 A (MASIAKOWSKI PIOTR J ;PROCTER & GAMBLE COMPAGNY (US); MORRIS JODI () 18 February 1999 (1999-02-18) claims ---	1
P,X	WO 99 26977 A (TSCHOPP JURG ;BIOGEN INC (US)) 3 June 1999 (1999-06-03)	1
P,A	the whole document ---	6,10,13
P,X	WO 99 04001 A (ZYMOGENETICS INC) 28 January 1999 (1999-01-28)	1
P,A	sequence ID no.2 claims ---	6,10,13
T	G. MATUTE-BELLO ET AL: "Soluble Fas Ligand induces epithelial cell apoptosis in humans with acute lung injury (ARDS)" JOURNAL OF IMMUNOLOGY., vol. 163, no. 4, 15 August 1999 (1999-08-15), pages 2217-2225, XP002137439 BALTIMORE US -----	

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 99/30734

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Although claim 13 is directed to a method of treatment of the human/animal body (rule 39.1 IV PCT), the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Claims 1, 6, 10, 13.

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1,6,10,13

mFLINT in the treatment of acute respiratory distress syndrome

2. Claims: 2,3,7,8,11,12,14,15

mFLINT in the treatment and/inhibition of ulcerative colitis.

3. Claims: 4,9

mFLINT for use in inhibiting ischemic injury during organ transplantation.

4. Claims: 5,16

mFLINT for use in organ preservation during transplantation.

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/30734

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0861850 A	02-09-1998	US 5885800 A	23-03-1999
		CA 2220852 A	03-08-1998
		JP 10215886 A	18-08-1998
WO 9830694 A	16-07-1998	AU 5815798 A	03-08-1998
		AU 6238698 A	03-08-1998
		CN 1247567 T	15-03-2000
		EP 1007659 A	14-06-2000
		EP 0990031 A	05-04-2000
		WO 9830693 A	16-07-1998
WO 9950413 A	07-10-1999	AU 3369199 A	18-10-1999
		WO 0037094 A	29-06-2000
WO 9914330 A	25-03-1999	AU 9497098 A	05-04-1999
		EP 1015587 A	05-07-2000
WO 9907738 A	18-02-1999	AU 8767698 A	01-03-1999
		EP 1019502 A	19-07-2000
WO 9926977 A	03-06-1999	AU 1535699 A	15-06-1999
		ZA 9810745 A	24-05-1999
WO 9904001 A	28-01-1999	AU 9013998 A	10-02-1999